

Matthew R. Fogg, Esq. DE Bar #4254
Keith E. Donovan, Esq. DE Bar #3044
MORRIS JAMES LLP
3205 Avenue North Blvd., Ste. 100
Wilmington, DE 19803
mfogg@morrisjames.com
kdonovan@morrisjames.com
(302) 655-8808 (Telephone)

Brandon L. Bogle, Esq. FL Bar #52624
Madison A. Mougey, Esq. FL Bar # 1071135
**LEVIN, PAPANTONIO, PROCTOR, BUCHANAN,
O'BRIEN, BARR & MOUGEY, P.A.**
316 South Baylen Street, Suite 600
Pensacola, Florida 32502
bbogle@levinlaw.com
mmougey@levinlaw.com
(850) 435-7043 (Telephone)

**UNITED STATES DISTRICT COURT
DISTRICT OF NEW JERSEY**

**IN RE: DUPIXENT (DUPILUMAB)
PRODUCTS LIABILITY LITIGATION**

Civil Action No. 26-md-3180

This Document Relates To:

ANNE GIBSON,

Plaintiff,

vs.

**SANOFI-AVENTIS U.S. LLC, GENZYME
CORPORATION, REGENERON
PHARMACEUTICALS, INC.**

Defendants.

COMPLAINT FOR DAMAGES

DEMAND FOR JURY TRIAL

COMPLAINT

COMES NOW Plaintiff, Anne Gibson, residing at 209 Grand County Road 662, Grand Lake, Colorado 80447, by and through undersigned counsel, and brings this action for damages

against Defendants Sanofi-Aventis U.S. LLC, Genzyme Corporation, and Regeneron Pharmaceuticals, Inc. Defendant Sanofi-Aventis U.S. LLC is a limited liability company with its principal place of business at 55 Corporate Drive, Bridgewater, NJ 08807. Defendant Genzyme Corporation is a corporation with its principal place of business at 450 Water Street, Cambridge, MA 02141. Defendant Regeneron Pharmaceuticals, Inc. is a corporation with its principal place of business at 777 Old Saw Mill Road, Tarrytown, NY 10591.

INTRODUCTION

1. This is an action for damages due to Plaintiff relating to Defendants' development, testing, manufacture, packaging, preparation, labeling, marketing, supply and/or sale of the dangerous and defective pharmaceutical product Dupixent®.

2. Defendants misrepresented that Dupixent was a safe and effective treatment for atopic dermatitis and other inflammatory health conditions when in fact the drug causes cutaneous t-cell lymphoma (CTCL), a form of non-Hodgkin's lymphoma, and/or triggers a reaction leading to severe and rapid clinical progression of otherwise subclinical and undiagnosed CTCL.

3. Defendants failed to warn physicians and the public about Dupixent's propensity to cause CTCL or rapidly exacerbate subclinical CTCL.

4. Consumers and physicians alike have been misled about the safety and efficacy of Dupixent, and as a result consumers, including Plaintiff, have developed CTCL.

PARTIES

5. Plaintiff, Anne Gibson is and was at all times relevant hereto, a resident of Grand County, Colorado.

6. Defendant, Sanofi-Aventis U.S. LLC, a wholly-owned subsidiary of Sanofi, is and was at all times relevant hereto a limited liability company organized and existing under the laws

of the State of Delaware, having its principal place of business at 55 Corporate Drive, Bridgewater, New Jersey 08807. Upon information and belief, Defendant, Sanofi-Aventis U.S. LLC has been substantively involved in, *inter alia*, the development and sale of Dupixent and the design, funding, conduct, authorship, publication and dissemination of scientific research and promotional messaging related to Dupixent and the indications for which Dupixent has received approval to treat.

7. Defendant, Genzyme Corporation, a wholly-owned subsidiary of Sanofi, is and was at all times relevant hereto a corporation organized and existing under the laws of the State of Massachusetts, having its principal place of business at 450 Water Street, Cambridge, Massachusetts 02142. Upon information and belief, Defendant, Genzyme Corporation has done business under various names and/or aliases, including “Sanofi Genzyme” and “Sanofi-Genzyme”, at certain times relevant to this action. Defendant, Genzyme Corporation’s headquarters is located less than one mile from Defendant, Regeneron Pharmaceuticals, Inc.’s Massachusetts office. Genzyme Corporation “focuses on developing specialty treatments for debilitating diseases that are often difficult to diagnose and treat, providing hope to patients and their families”.¹ At the time of launch, Dupixent was considered an “[i]mportant new addition to Sanofi Genzyme [Genzyme Corporation] specialty care business.”² Defendant, Genzyme Corporation has been substantively involved in, *inter alia*, the coordination and funding of third parties to design, conduct, author, publish and disseminate scientific research and promotional messaging related to Dupixent and the indications for which Dupixent has received approval to treat. Prior to Dupixent’s launch,

¹ Regeneron Pharm., Inc. & Sanofi, Regeneron and Sanofi Announce Positive Dupixent® (Dupilumab) Phase III Data, Regeneron (Oct. 27, 2024), <https://investor.regeneron.com/news-releases/news-release-details/regeneron-and-sanofi-announce-positive-dupixentr-dupilumab-phase>

² Sanofi, *Dupixent FDA Approval Investor Call* (Mar. 28, 2017), https://www.sanofi.com/assets/dotcom/content-app/events/investor-presentation/2017/Dupixent_FDA_Approval_Investor_call_FINAL.pdf (last visited Nov. 25, 2025).

Defendant Genzyme Corporation worked hand in hand with Defendants Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC in commercializing Dupixent.³

8. Defendant, Regeneron Pharmaceuticals, Inc. is and was at all times relevant hereto a corporation organized under the laws of the state of New York with its principal place of business at 777 Old Saw Mill River Road, Tarrytown, New York 10591. Regeneron Pharmaceuticals, Inc. lists nine locations in North America, including Cambridge, Massachusetts. Defendant, Regeneron Pharmaceuticals, Inc. is the current official sponsor of the Biologics License Application for Dupixent in the United States, and thus maintains significant responsibility and control over the drug and all activities and materials relating thereto. Defendant, Regeneron Pharmaceuticals, Inc. has been substantively involved in the development and sale of Dupixent and the design, funding, conduct, authorship, publication and dissemination of scientific research and promotional messaging related to Dupixent and the indications for which Dupixent has received approval to treat.

9. Defendants, Regeneron Pharmaceuticals, Inc.; Sanofi-Aventis U.S. LLC; and Genzyme Corporation, shall hereinafter be referred to collectively as “Defendants”.

10. Defendants, directly and/or through third parties acting upon Defendants’ directives, were jointly engaged in the business of designing, developing, manufacturing, testing, packaging, promoting, marketing, distributing, labeling, and/or selling Dupixent and controlling the Biologics License Application for Dupixent and other intellectual property associated with Dupixent.

11. At all times relevant hereto, Defendants include and have included any and all

³ Regeneron Pharm., Inc. & Sanofi, Regeneron and Sanofi Announce Marketing Authorization Application for Dupixent® (dupilumab) Accepted for Review by the EMA (Dec. 8, 2016), <https://investor.regeneron.com/news-releases/news-release-details/regeneron-and-sanofi-announce-marketing-authorization>

parent companies, subsidiaries, affiliates, divisions, franchises, partners, joint venturers, and organizational units of any kind, their predecessors, successors and assigns and their officers, directors, employees, agents, representatives and any and all other persons acting on their behalf.

12. At all times relevant hereto, Defendants were engaged in the business of developing, designing, licensing, manufacturing, distributing, selling, marketing, and or introducing into interstate commerce throughout the United States, and in the state of Colorado, either directly or indirectly, through third-parties, subsidiaries and/or related entities, the pharmaceutical product Dupixent.

13. Consistent with Plaintiff's allegations herein, in *Immunex Corp. v. Sanofi*, No. 2:17-cv-02613, 2017 (C.D. Cal. Aug. 16, 2017), Defendants expressly admitted in their amended answer, (Doc. No. 140) the following:

- a. Defendants, collectively, manufacture, use, and sell Dupixent. Am. Answer ¶ 1, Doc. 104.
- b. Defendants Genzyme Corporation and Regeneron Pharmaceuticals, Inc. have marketed and sold Dupixent. Am. Answer ¶ 46, Doc. 104.
- c. Defendant Genzyme Corporation prepared a U.S.-based salesforce to offer for sale and to sell Dupixent in the United States. Am. Answer ¶ 47, Doc. 104.

JURISDICTION & VENUE

14. The jurisdiction of this Court over the subject matter of this action is predicated on 28 U.S.C. § 1332. The amount in controversy exceeds \$75,000.00, exclusive of interest and costs and complete diversity of citizenship exists between the parties.

15. Venue in this Court is proper pursuant to 28 U.S.C § 1391 in that a substantial part of the events or omissions giving rise to the claims asserted herein occurred in this District, and Defendants are subject to personal jurisdiction in this District.

GENERAL ALLEGATIONS

16. This action is for damages brought on behalf of Plaintiff, Anne Gibson, who was prescribed and supplied with and who has taken the prescription drug Dupixent, as tested, studied, researched, evaluated, endorsed, designed, formulated, compounded, manufactured, produced, processed, assembled, inspected, distributed, marketed, labeled, promoted, packaged, advertised for sale, or otherwise placed in the stream of commerce by Defendants.

17. Plaintiff used Dupixent to treat atopic dermatitis from March 2021 to February 2023. Thereafter Plaintiff was diagnosed with CTCL.

18. Plaintiff, Anne Gibson, brings this action against Defendants to recover damages for the injuries suffered as a result of her use of Dupixent and to recover for her individual economic and non-economic damages which she sustained as a result therefrom.

19. Defendants' wrongful acts, omissions, and fraudulent misrepresentations caused Plaintiff's injuries and damages.

20. At all times relevant, Defendants were engaged in the business of researching, licensing, designing, formulating, compounding, testing, manufacturing, producing, processing, assembling, inspecting, distributing, marketing, labeling, promoting, packaging and/or advertising for sale the prescription drug Dupixent for use by physicians in treating their patients, including Plaintiff.

21. At all times relevant hereto, Defendants were authorized to do business within Plaintiff's state of residence and did conduct such business.

22. At times relevant hereto, the officers and directors of Defendants participated in, authorized, and directed the production and promotion of Dupixent when they knew, or with the exercise of reasonable care should have known, of the hazards and dangerous propensities of

Dupixent and thereby actively participated in the tortious conduct which resulted in the injuries suffered by Plaintiff discussed herein.

FACTS COMMON TO ALL COUNTS

A. Dupixent

23. Dupixent® (dupilumab) is a therapeutic biologic used in the treatment of infants, children and adults with inflammatory health conditions in the United States and worldwide.

24. Dupixent is a human monoclonal immunoglobulin G4 antibody that binds to the interleukin (IL)-4 receptor alpha subunit shared by IL-4 and IL-13 to prevent their signaling. IL-4 and IL-13 signaling produces an inflammatory cascade that has been implicated in a number of immune-mediated inflammatory conditions, including atopic dermatitis, chronic rhinosinusitis with nasal polyposis, eosinophilic esophagitis, prurigo nodularis, chronic obstructive pulmonary disease, chronic spontaneous urticaria and bullous pemphigoid. Despite this, the mechanism of action through which Dupixent exerts a therapeutic effect in patients with these conditions has not been definitively established.

25. Dupixent is supplied by Defendants in 200 mg and 300 mg single-dose prefilled syringes and single-dose prefilled pens. Dupixent is self-administered by patients or by caregivers through a subcutaneous injection in 1-, 2- or 4-week intervals depending on age, weight and therapeutic indication.

B. Defendants' Collaboration Agreement

26. In September 2003, Defendant, Regeneron Pharmaceuticals, Inc. entered into a Collaboration Agreement with Aventis Pharmaceuticals Inc. to co-develop an oncology medication it had invented.

27. In or about December 2004, Aventis Pharmaceuticals Inc. merged with and into

Sanofi, at the time operating under the name Sanofi-Aventis, and formerly known as Sanofi-Synthelabo. Upon information and belief, Sanofi, at the time operating under the name Sanofi-Aventis, became the successor in interest to Aventis Pharmaceuticals Inc. in all agreements to which Aventis Pharmaceuticals Inc. was a party and acquired all rights formerly held by Aventis Pharmaceuticals Inc. in its Collaboration Agreement with Defendant, Regeneron Pharmaceuticals, Inc.

28. This Collaboration Agreement was again amended in January 2005, December 2005 and January 2006, at which point Defendant, Sanofi-Aventis U.S., LLC was named the successor in interest to Aventis Pharmaceuticals Inc. in the Collaboration Agreement with Defendant, Regeneron Pharmaceuticals, Inc.

29. In November 2007, Defendant, Regeneron Pharmaceuticals, Inc. entered into a new License and Collaboration Agreement with Sanofi, via its subsidiaries and/or affiliate entities. Under this License and Collaboration Agreement, Sanofi agreed to provide research funding in exchange for exclusive rights to co-develop and co-commercialize any new biopharmaceuticals discovered by Defendant, Regeneron Pharmaceuticals, Inc. In this agreement, Defendant, Sanofi-Aventis U.S. LLC was again identified as the successor in interest to Aventis Pharmaceuticals Inc. in the original September 2003 Collaboration Agreement and all amendments thereto. Defendants contemporaneously entered into a Discovery and Preclinical Development Agreement under which Defendant, Regeneron Pharmaceuticals, Inc. would discover and create products for joint development and commercialization with Sanofi and its subsidiaries and/or affiliates. Such products would be considered a “Licensed Product” under the terms of the License and Collaboration Agreement.

30. In November 2009, Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi, via

its subsidiaries and/or affiliate entities, expanded the scope and terms of their collaboration in an Amended and Restated License and Collaboration Agreement and Amended and Restated Discovery and Preclinical Development Agreement. The Amended and Restated License and Collaboration Agreement currently remains in effect, with subsequent amendments in May 2013, July 2015, April 2020 and September 2021, while the Amended Discovery Agreement expired in 2017.

31. Upon information and belief, as of January 2014, Defendant, Sanofi-Aventis U.S. LLC was again named successor in interest to Aventis Pharmaceuticals Inc. with respect to the original September 2003 Collaboration Agreement and all amendments thereto, as well as the First Amendment to the Amended and Restated License and Collaboration Agreement dated May 2013.

32. One of the drugs Defendants agreed to co-commercialize under the Amended and Restated License and Collaboration Agreement was Dupixent. Sanofi via its subsidiary and/or affiliate entities provided upfront funding to cover 100% of the costs incurred by Defendant, Regeneron Pharmaceuticals, Inc. in the development of Dupixent and other Licensed Products under the Amended and Restated License and Collaboration Agreement, including an initial non-refundable sum of \$85 million USD.

33. Sanofi, individually and/or through its subsidiaries, is considered the lead party with respect to the commercialization of Dupixent, and Defendant, Regeneron Pharmaceuticals, Inc. exercised its co-promotion rights under the Restated License and Collaboration Agreement for Dupixent in the United States.

34. Sanofi, individually and/or through its subsidiary, Defendant, Sanofi-Aventis U.S. LLC, continues to supply funding for ongoing Dupixent development research being conducted by Defendant, Regeneron Pharmaceuticals, Inc.

35. Upon information and belief, Defendants, Regeneron Pharmaceuticals, Inc., Sanofi-Aventis U.S. LLC, and Genzyme Corporation, both independently and collaboratively, are substantively involved in performing commercialization activities for Dupixent in the United States, including but not limited to manufacturing, producing, processing, assembling, inspecting, distributing, marketing, labeling, promoting, packaging and/or advertising, along with support activities associated with each of those functions, both directly and through third parties acting on their behalf.

36. Under the terms of the Amended and Restated License and Collaboration Agreement, Defendants have a reciprocal duty to regularly share information relating to the commercialization of Dupixent, to include anticipated launch dates, key market metrics, market research and sales, as well as to promptly notify the other party of any major market developments.

37. Sanofi, the parent company of Defendants, Sanofi-Aventis U.S. LLC and Genzyme Corporation, recognizes all sales of Dupixent. Profits and losses arising from commercial operations in the United States, including those associated with Dupixent, are split equally (50/50) between Sanofi and its subsidiaries and Defendant, Regeneron Pharmaceuticals, Inc. Profit-sharing between Defendants for sales of Dupixent outside of the United States is done based on a sliding scale. Defendant, Regeneron Pharmaceuticals, Inc. also receives or has at times received milestone payments from Sanofi based on sales of Dupixent and other Licensed Products in markets outside of the United States.

38. Upon information and belief, Sanofi serves as the “Lead Regulatory Party”, as defined in the Amended and Restated License and Collaboration Agreement, responsible for managing pharmacovigilance and product complaints and for formulating and implementing any related strategies for Dupixent in the United States. Further upon information and belief,

Defendant, Sanofi-Aventis U.S. LLC handles all processing and submissions of adverse event reports and product complaints for Dupixent in the United States on behalf of Sanofi. Despite this, Defendants have joint responsibility under the terms of the Amended and Restated License and Collaboration Agreement to collaborate in fulfilling all regulatory requirements concerning pharmacovigilance and risk management plans and product complaint reporting associated with Dupixent in the United States. Further to this joint responsibility, upon information and belief, Defendants executed a Safety Data Exchange Agreement setting forth the specific procedures to be used to coordinate the investigation and exchange of reports of adverse events and complaints associated with Dupixent to ensure timely communication to regulatory authorities and compliance with applicable laws as prescribed in the Amended and Restated License and Collaboration Agreement.

C. Development of Dupixent

39. Upon information and belief, the human monoclonal antibody and active ingredient in Dupixent, dupilumab, formerly known under the common names REGN668 and SAR231893, was originally discovered by Defendant, Regeneron Pharmaceuticals, Inc. during the mid- to late-2000s. Defendant, Regeneron Pharmaceuticals, Inc. utilized its proprietary VelocImmune® technology, a genetically-engineered mouse platform endowed with a genetically-humanized immune system, in order to invent dupilumab.

40. Defendant, Regeneron Pharmaceuticals, Inc. has been assigned at least 7 United States patents relating to Dupixent, including US 7,608,693 B2 (October 2009), US 8,735,095 B2 (May 2014), US 8,945,559 B2 (February 2015), US 9,238,692 B2 (January 2016), US 10,435,473 B2 (October 2019), US 11,059,896 B2 (July 2021) and US 11,926,670 (March 2024).

41. The first clinical development program for Dupixent in the treatment of patients

with atopic dermatitis was regulated under Investigational New Drug (“IND”) Application number 107969 in the United States. Defendant, Regeneron Pharmaceuticals, Inc. submitted IND number 107969 to the United States Food and Drug Administration (“FDA”) on July 12, 2010 requesting permission to study Dupixent in human subjects. The FDA deemed the first Phase I clinical study to assess the safety and tolerability of Dupixent in adult patients with atopic dermatitis “Safe to Proceed” on August 21, 2010.

42. During clinical development, the FDA granted Dupixent a Breakthrough designation, Priority Review and Rolling Review, regulatory programs which are each intended to expedite development and approval of new therapeutics. Defendant, Regeneron Pharmaceuticals, Inc., completed its rolling submission of Biologics License Application (“BLA”) number 761055 for Dupixent to FDA on July 28, 2016. FDA received the BLA on July 29, 2016.

43. Defendants officially received FDA approval to market Dupixent in the United States on March 28, 2017 for the treatment of adult patients with moderate to severe atopic dermatitis whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable.

44. The original approval of Dupixent was based on the results of three Phase III randomized controlled trials, SOLO 1, SOLO 2, and CHRONOS, which used a clinical scale known as the Investigator’s Global Assessment (IGA) to measure changes in atopic dermatitis disease severity in response to treatment. Each of these trials demonstrated superiority of Dupixent over placebo or placebo plus topical corticosteroids in the primary endpoint, change from baseline to Week 16 in the proportion of subjects achieving an IGA score of 0 (clear skin) or 1 (almost clear skin), with at least a 2-point improvement in IGA score.

45. Following its approval in March 2017 as a second-line treatment for adult patients

with atopic dermatitis in the United States, Defendants proceeded to submit several supplemental BLAs to the FDA seeking to market Dupixent for the treatment of additional health conditions and expand the indicated patient population to include nearly all age groups.

46. In October 2018, Dupixent was approved as an add-on maintenance treatment for patients aged 12 years and older with moderate-to-severe asthma characterized by an eosinophilic phenotype or with oral corticosteroid dependent asthma.

47. In March 2019, Dupixent's indicated atopic dermatitis population was expanded to include patients aged 12 years and older.

48. In June 2019, Dupixent was approved as an add-on maintenance treatment in adult patients with inadequately controlled chronic rhinosinusitis with nasal polyps.

49. In May 2020, Dupixent's indicated atopic dermatitis population was expanded to include patients aged 6 years and older.

50. In October 2021, Dupixent's indicated asthma population was expanded to include patients aged 6 years and older.

51. In May 2022, Dupixent was approved for the treatment of patients aged 12 years and older with eosinophilic esophagitis.

52. In June 2022, Dupixent's indicated atopic dermatitis population was expanded to include patients aged 6 months and older.

53. In September 2022, Dupixent was approved for the treatment of adult patients with prurigo nodularis.

54. In January 2024, Dupixent's indicated eosinophilic esophagitis population was expanded to include patients aged 1 year and older.

55. In September 2024, Dupixent's indicated chronic rhinosinusitis with nasal polyps

population was expanded to include patients aged 12 years and older. Dupixent was also approved as an add-on maintenance treatment of adult patients with inadequately controlled chronic obstructive pulmonary disease and an eosinophilic phenotype in September 2024.

56. In April 2025, Dupixent was approved for the treatment of patients aged 12 years and older with chronic spontaneous urticaria who remain symptomatic despite H1 antihistamine treatment.

57. Finally, in June 2025, Dupixent was approved for the treatment of adult patients with bullous pemphigoid.

58. Defendants continue to co-develop Dupixent under INDs in the treatment of chronic pruritus of unknown origin, eosinophilic gastritis with or without eosinophilic duodenitis, lichen simplex chronicus and ulcerative colitis.

D. Defendants' Marketing of Dupixent

59. Defendants have utilized multiple mediums and mechanisms to aggressively promote Dupixent as a necessary, life-changing therapy for patients suffering from atopic dermatitis and other inflammatory health conditions in order to drive prescriptions for Dupixent. Much of Defendants' approach to boosting Dupixent uptake has involved expanding the market for its use by convincing individuals with subtle symptoms, or even no symptoms, that they actually have a serious, undiagnosed inflammatory health condition, and penetrating the existing market by convincing patients that their current treatment regimen is inadequate to control their disease and/or has an unacceptable safety profile.

60. Defendants have infused billions of dollars into persuasive direct-to-consumer advertising depicting Dupixent as being highly safe and effective and capable of greatly improving the quality of patients' lives.

61. In 2024, Defendants spent an estimated \$484.1 million on advertising for Dupixent across all media types. This spend included \$276 million on 17 different direct-to-consumer television advertisements: 8 for atopic dermatitis, 8 for asthma and 1 for gastrointestinal conditions. Defendants' annual advertising expenditures for Dupixent were the third highest among all marketed drugs in 2024.⁴

62. In 2023, Defendants spent an estimated \$502 million on advertising for Dupixent across all media types. This spend included \$315.3 million on 12 different direct-to-consumer television advertisements: 8 for atopic dermatitis and 4 for asthma. Defendants' annual advertising expenditures for Dupixent were the second highest among all marketed drugs in 2023.⁵

63. In 2022, Defendants spent an estimated \$491 million on advertising for Dupixent across all media types. This spend included \$305.9 million on direct-to-consumer television advertisements. Defendants' annual advertising expenditures for Dupixent were higher than any other marketed drugs in 2022.⁶

64. In 2021, Defendants spent an estimated \$523.9 million on advertising for Dupixent across all media types. This spend included \$287.6 million on direct-to-consumer television

⁴ Andrea Park, *AbbVie Pulls Off a Hat Trick with 3rd Straight Year as Top TV Drug Ad Spender, Buoyed by Skyrizi and Rinvoq Spots*, Fierce Pharma (Jan. 22, 2025), <https://www.fiercepharma.com/marketing/abbvie-pulls-hat-trick-3rd-straight-year-top-tv-drug-ad-spender-buoyed-skyrizi-and-rinvoq> (last visited Dec. 9, 2025); Andrea Park, Nick Paul Taylor & Zoey Becker, *The Top 10 Pharma Drug Ad Spenders of 2024*, Fierce Pharma (June 30, 2025), <https://www.fiercepharma.com/marketing/top-10-pharma-drug-ad-spenders-2024> (last visited Dec. 9, 2025).

⁵ Ben Adams, Andrea Park & Nick Paul Taylor, *AbbVie Makes Clean Sweep—Skyrizi and Rinvoq Top TV Drug-Ad Spender for 2023*, Fierce Pharma (Jan. 9, 2024), <https://www.fiercepharma.com/marketing/abbvie-makes-clean-sweep-skyrizi-and-rinvoq-top-tv-drug-ad-spender-2023> (last visited Dec. 9, 2025); Ben Adams, Andrea Park & Nick Paul Taylor, *The Top 10 Pharma Drug Ad Spenders for 2023*, Fierce Pharma (June 3, 2024), <https://www.fiercepharma.com/marketing/top-10-pharma-drug-ad-spenders-2023> (last visited Dec. 9, 2025); Park, supra note 5, <https://www.fiercepharma.com/marketing/abbvie-pulls-hat-trick-3rd-straight-year-top-tv-drug-ad-spender-buoyed-skyrizi-and-rinvoq>; Andrea Park, Nick Paul Taylor & Zoey Becker, *The Top 10 Pharma Drug Ad Spenders of 2024*, Fierce Pharma (June 30, 2025), <https://www.fiercepharma.com/marketing/top-10-pharma-drug-ad-spenders-2024> (last visited Dec. 9, 2025).

⁶ Ben Adams, *The Top 10 Pharma Drug-Brand Ad Spenders for 2022*, Fierce Pharma (May 1, 2023), <https://www.fiercepharma.com/special-reports/top-10-pharma-drug-brand-ad-spenders-2022> (last visited Dec. 9, 2025).

advertisements. Defendants' annual advertising expenditures for Dupixent were higher than any other marketed drugs in 2021.⁷

65. In 2020, Defendants spent an estimated \$409.8 million on advertising for Dupixent across all media types. This spend included millions on 7 different direct-to-consumer television advertisements: 4 for atopic dermatitis and 3 for asthma. Defendants' annual advertising expenditures for Dupixent were the second highest among all marketed drugs in 2020.⁸

66. In 2019, Defendants spent an estimated \$199.4 million on their first full year of advertising for Dupixent across all media types. This spend included \$76 million on direct-to-consumer television advertisements. Defendants' annual advertising expenditures for Dupixent were the third highest among all marketed drugs in 2019.⁹

67. In 2018, Defendants spent an estimated \$104.2 million on direct-to-consumer advertising.¹⁰

68. Defendants' provocative messaging typically bombards consumers with glossy imagery and catchy phrases like "Du-More", "Better Days" and "This is Better" that imply patients will lead a happier, more productive and fulfilling life if they use Dupixent. According to a spokesperson for Defendant, Regeneron Pharmaceuticals, Inc., one 60-second Dupixent commercial that aired throughout 2024 made patients "not only want to talk to their doctors but

⁷ *Id.*

⁸ Beth Snyder Bulik, *The Top 10 Ad Spenders in Big Pharma for 2020*, Fierce Pharma (Apr. 19, 2021), <https://www.fiercepharma.com/special-report/top-10-ad-spenders-big-pharma-for-2020> (last visited Dec. 9, 2025).

⁹ Beth Snyder Bulik, *Pharma's 2019 TV Spending Ticks Up — Barely — Thanks to Big Humira Growth*, Fierce Pharma (Jan. 14, 2020), <https://www.fiercepharma.com/marketing/pharma-tv-ad-spending-for-2019-finishes-flat-compared-to-previous-year> (last visited Dec. 9, 2025); Beth Snyder Bulik, *The Top 10 Ad Spenders in Big Pharma for 2019*, Fierce Pharma (Feb. 19, 2020), <https://www.fiercepharma.com/special-report/top-10-advertisers-big-pharma-for-2019> (last visited Dec. 9, 2025).

¹⁰ U.S. Gov't Accountability Off., GAO-21-380, *Prescription Drugs: Medicare Spending on Drugs with Direct-to-Consumer Advertising: Report to the Committee on the Judiciary, U.S. Senate* (May 2021), <https://www.gao.gov/assets/gao-21-380.pdf> (last visited Dec. 9, 2025).

also to rewatch the ad”.¹¹

69. To ensure ready access to Dupixent and prevent potential sales lapses from treatment interruptions or omissions, Defendants launched Dupixent MyWay, a support program that helps patients “get access to DUPIXENT and stay on track”.¹² Defendants paired Dupixent MyWay with the Dupixent MyWay Ambassador program which uses real world patients to endorse and further disseminate their messaging. Defendants also created and launched smartphone apps, EZTtrack for Atopic Dermatitis and the Dupixent MyWay Patient App, to help patients access Dupixent “as quickly as possible” and facilitate “more productive conversations” with doctors that are more likely to lead to a prescription for Dupixent.^{13,14}

70. In conjunction with branded advertising, Defendants have amplified public awareness of the conditions for which Dupixent is indicated in order to increase rates of their diagnosis and further drive prescriptions of Dupixent. Defendants accomplished this by using unbranded disease state awareness websites and funding and conducting large-scale unbranded disease state awareness campaigns, scientific publications and continuing medical education programs. With these multifaceted unbranded campaigns, which Defendants designed to track branded marketing campaigns, Defendants began seeding the market for Dupixent well before it received FDA approval for marketing in the United States.

71. Defendant, Regeneron Pharmaceuticals, Inc. initially registered the disease state awareness website UnderstandAD.com on March 23, 2016 and continues to own and operate it in

¹¹ Andrea Park, Nick Paul Taylor & Zoey Becker, *The Top 10 Pharma Drug Ad Spenders of 2024*, Fierce Pharma (June 30, 2025), <https://www.fiercepharma.com/marketing/top-10-pharma-drug-ad-spenders-2024> (last visited Dec. 9, 2025).

¹² DUPIXENT MyWay, *Patient Support Program*, dupixent.com, <https://www.dupixent.com/support-savings/dupixent-my-way> (last visited Dec. 9, 2025).

¹³ Apple Inc., *MyWay Patient Support*, Apple App Store, <https://apps.apple.com/us/app/myway-patient-support/id6443848439> (last visited Dec. 9, 2025).

¹⁴ Apptopia, About — *MyWay Patient Support (iOS)*, Apptopia, <https://apptopia.com/ios/app/1501488844/about> (last visited Dec. 9, 2025).

conjunction with Defendant, Sanofi-Aventis U.S. LLC and the National Eczema Association. Also on March 23, 2016, Sanofi, under its former name Sanofi-Synthelabo, initially registered the disease state awareness website EczemaExposed.com which is currently hosted by Defendant, Sanofi-Aventis U.S. LLC. Defendants use these disease state awareness websites to provide seemingly unbiased educational information geared towards highlighting the potential burdens of atopic dermatitis on patients, helping patients identify signs and symptoms of undiagnosed atopic dermatitis and suggesting to patients that their condition is more uncontrolled than they might believe, while also supplying guidance to steer productive communications with physicians regarding available treatment options. Upon information and belief, Defendants registered these web domains approximately one week before making their first rolling submission of the Dupixent BLA to the FDA.

72. In Defendant Regeneron Pharmaceuticals, Inc.'s, November 2017 Earnings Call, Defendants acknowledged their goal to increase rates of diagnosis and further drive prescriptions of Dupixent. Robert J. Terifay, Regeneron's Executive Vice President, Commercial, stated "we anticipate is that we will see other prescribers as patients become aware through our direct-to-consumer campaign on the severity of the condition and the need to seek out a dermatologist, and that there could be an increased urgency to treat over time".¹⁵

73. During this call, Defendants also represented prescriptions are not driven from physicians but rather ". . . expansion into moderate AD. . . helped by our direct-to-consumer campaign".¹⁶

74. Defendants' vast unbranded marketing campaigns for other indicated conditions

¹⁵ Regeneron Pharm., Inc., Q3 2017 Earnings Call Transcript, ROIC.AI (Nov. 8, 2017), <https://www.roic.ai/quote/REGN/transcripts/2017/3>

¹⁶ *Id.*

have followed suit. Another disease state awareness website targeting asthma and chronic rhinosinusitis with nasal polyps, TheNextBreath.com, was originally registered by Sanofi, under its former name Sanofi-Synthelabo, and is currently maintained by Defendant, Regeneron Pharmaceuticals, Inc. and Sanofi, individually and/or through its subsidiaries. Through this website Defendants aim to “bring more awareness to severe asthma” and “empower people to take action to strive for better asthma control”, while claiming asthma “often goes unrecognized”. Additionally, Defendants cite statistics such as “[o]ver 80% of people overestimate how well controlled their asthma is” and nearly half of asthma patients “have symptoms that are not well controlled”, as a means to convince patients that believe they have “good” or “very good” control over their asthma symptoms that they are actually wrong.

75. Another disease state awareness website, Type2Inflammation.com, was originally registered by Sanofi, under its former name Sanofi-Synthelabo, in February 2019 and is currently maintained by Defendant, Sanofi-Aventis U.S. LLC on behalf of Defendants. Defendants utilize this unbranded website to drive awareness and treatment of several conditions for which Dupixent has been approved, including eosinophilic esophagitis, chronic spontaneous urticaria, prurigo nodularis, atopic dermatitis, asthma, chronic rhinosinusitis with nasal polyposis and chronic obstructive pulmonary disease.

76. Finally, Defendants own and maintain two disease state awareness websites dedicated to driving awareness and treatment of chronic obstructive pulmonary disease: ThinkCOPDInflammation.com, initially registered in March 2023 by Sanofi, under its former name Sanofi-Synthelabo, and currently maintained by Defendant, Regeneron Pharmaceuticals, Inc. and Sanofi, individually and/or through its subsidiaries; and LiveWithCOPD.com, initially registered in December 2023 by Sanofi, under its former name Sanofi-Synthelabo, and currently

maintained by Defendant, Sanofi-Aventis U.S. LLC on behalf of Defendants. Upon information and belief, Defendants registered the web domain ThinkCOPDInflammation.com approximately 9 months before submitting the supplemental BLA for Dupixent in the treatment of COPD to the FDA, and Defendants registered the web domain LiveWithCOPD.com approximately 2 weeks before submitting the supplemental BLA for Dupixent in the treatment of COPD to the FDA.

77. On July 12, 2016, Defendants, Regeneron Pharmaceuticals, Inc., Sanofi-Aventis U.S. LLC, and Genzyme Corporation launched the “Understand AD” awareness campaign which was “focused on educating people about moderate-to-severe atopic dermatitis”.¹⁷ Defendants highlighted extreme examples of symptoms to depict how uncontrolled atopic dermatitis can impact patients’ lives and utilized celebrity chef Elizabeth Falkner as a spokesperson to expand the reach and influence of this initiative. In a press release about the Understand AD campaign, Defendants specifically signaled “there is still a need for additional treatment options for atopic dermatitis”, while featuring quotes from leaders at the National Eczema Association and Dermatology Nurses Association regarding “raising the level of awareness” of atopic dermatitis, “advocating for better care and treatments” and “driv[ing] the dialogue that atopic dermatitis is more than skin deep”. Upon information and belief, Defendants launched this disease state awareness campaign approximately 2 weeks before making their final rolling submission of the Dupixent BLA to the FDA.

78. On October 11, 2017, Defendants, Regeneron Pharmaceuticals, Inc., Sanofi-Aventis U.S. LLC, and Genzyme Corporation launched the campaign “Understand AD: A Day in the Life” using Peter Moffat, writer and executive producer of the HBO show “The Night Of” to

¹⁷ Regeneron Pharmaceuticals, Inc., *Award-Winning Chef Elizabeth Falkner Reveals Her Struggle with Atopic Dermatitis to Highlight the Physical and Psychological Impact of the Disease* (news release), Regeneron Investor Relations (July 12, 2016), <https://investor.regeneron.com/news-releases/news-release-details/award-winning-chef-elizabeth-falkner-reveals-her-struggle-atopic> (last visited Dec. 9, 2025).

help push their unbranded disease awareness messaging for the stated purpose of “Driving Empathy and Understanding” of atopic dermatitis.¹⁸ In the spirit of disease mongering, Defendants represented that this program would serve to “transport[] people into what a typical, often unremitting and painful day is really like for people living with uncontrolled moderate-to-severe atopic dermatitis”. Taking care not to explicitly reference Dupixent, Defendants also casually highlighted that atopic dermatitis patients often “continue to experience debilitating symptoms despite available topical and systemic steroid treatment options, causing their disease to remain uncontrolled” in the press release announcing this campaign.

79. Then in September 2019 Defendants expanded the Understand AD program into its next phase, The Understand AD Squad video series, which was “targeted to adolescents with AD and their caregivers”.¹⁹ In the press release announcing the Understand AD Squad video series, Defendants exaggerated the rates and typical burdens of uncontrolled atopic dermatitis, suggesting “400,000 adolescents in the U.S. with uncontrolled moderate-to-severe atopic dermatitis” experience “debilitating, burdensome disease symptoms including intense, persistent itching, skin lesions and skin dryness, cracking, redness, crusting and oozing”.

80. The next step of Defendants’ calculated marketing efforts involved using published scientific literature to promote perceptions that patients suffering from atopic dermatitis are not being diagnosed or, if they have already received a diagnosis of atopic dermatitis, they are not being sufficiently treated.

¹⁸ Regeneron Pharmaceuticals, Inc., *Internationally Acclaimed Writer and Executive Producer of HBO’s “The Night Of” Partners With Regeneron to Release “Understand AD: A Day in the Life”* (news release), Regeneron Investor Relations (Oct. 11, 2017), <https://investor.regeneron.com/static-files/4e57c3b2-9384-451d-b508-5e41052cb738> (last visited Dec. 9, 2025).

¹⁹ Regeneron Pharmaceuticals, Inc., *Internationally Acclaimed Writer and Executive Producer of HBO’s “The Night Of” Partners With Regeneron to Release “Understand AD: A Day in the Life”* (news release), Regeneron Investor Relations (Oct. 11, 2017), <https://investor.regeneron.com/static-files/4e57c3b2-9384-451d-b508-5e41052cb738> (last visited Dec. 9, 2025).

81. Defendants wasted no time in strategizing Dupixent sales. Regeneron's Executive Vice President, Commercial, Robert J. Terifay, stated the month before FDA approval: "Sanofi Genzyme [Genzyme Corporation] and Regeneron have fully hired and trained our field teams".²⁰

82. In April 2017, the month following FDA approval of Dupixent for marketing in the United States, Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC funded and published a cross-sectional survey study via several of their employees to prompt increases in atopic dermatitis diagnoses and prescriptions of atopic dermatitis therapies.²¹ Published in American Journal of Clinical Dermatology, Defendants' study reported that awareness and use of clinical scales to measure atopic dermatitis severity was low, "suggesting a need for greater education of physicians regarding the availability and use of such measures for the assessment of AD". Despite nearly 70% concordance between patients and physicians on symptom severity, Defendants' employees concluded their results "show[ed] a discordance between patient- and physician-reported AD severity" and suggested "physicians may have a similar misunderstanding of the patient's perspective regardless of specialty". It was also the opinion of the authors that "clinical assessment may be less than optimal" and that "physicians may not be having meaningful conversations with patients about [quality of life] or the impact of AD on the daily life of the patient. The patient impact likely represents an important indicator of a need for treatment". Finally, the authors advised that it is important to include the patient perspective when making management decisions.

83. Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC then

²⁰ Regeneron Pharm., Inc., Q4 2016 Earnings Call Transcript, ROIC.AI (Feb. 23, 2017), <https://www.roic.ai/quote/REGN/transcripts/2016-year/4-quarter> (last visited Nov. 25, 2025).

²¹ Wei W, Anderson P, Gadkari A, et al. Discordance Between Physician- and Patient-Reported Disease Severity in Adults with Atopic Dermatitis: A US Cross-Sectional Survey. Am J Clin Dermatol. 2017;18(6):825-835. doi:10.1007/s40257-017-0284-y

funded and published a pair of studies in November 2017 and July 2018 in order to create and push a narrative that a majority of patients with atopic dermatitis continue to experience inadequate control of their symptoms despite treatment with therapies other than Dupixent, in addition to suffering from impaired quality of life and higher rates of anxiety, depression, work impairment, sleep impairment and itching symptoms.^{22,23}

84. Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC further endeavored to invent their own diagnostic tool for self-administration by patients. Defendants designed their 6-item patient symptom questionnaire, the Atopic Dermatitis Control Tool (ADCT), to have much greater sensitivity than existing tools, and as a result, its use by patients would serve to artificially increase rates of atopic dermatitis diagnoses. Defendants funded and published a pair of studies in November 2019 and December 2019 in order to introduce the ADCT.^{24,25} The ADCT tool has a dedicated website (adcontroltool.com) owned by Sanofi and maintained by Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC, and is currently available for patients to use on Defendants' unbranded website EczemaExposed.com. Defendants created similar questionnaires to use with and increase diagnoses of atopic dermatitis in children aged 6 to 11 years, the Worst Itch Scale, and children and infants aged 6 months to less than 6 years, the

²² Wei W, Anderson P, Gadkari A, et al. Extent and consequences of inadequate disease control among adults with a history of moderate to severe atopic dermatitis. *J Dermatol*. 2018;45(2):150-157. doi:10.1111/1346-8138.14116

²³ Simpson EL, Guttman-Yassky E, Margolis DJ, et al. Association of Inadequately Controlled Disease and Disease Severity With Patient-Reported Disease Burden in Adults With Atopic Dermatitis. *JAMA Dermatol*. 2018;154(8):903-912. doi:10.1001/jamadermatol.2018.1572

²⁴ Simpson E, Eckert L, Gadkari A, et al. Validation of the Atopic Dermatitis Control Tool (ADCT®) using a longitudinal survey of biologic-treated patients with atopic dermatitis. *BMC Dermatol*. 2019;19(1):15. Published 2019 Nov 6. doi:10.1186/s12895-019-0095-3

²⁵ Pariser DM, Simpson EL, Gadkari A, et al. Evaluating patient-perceived control of atopic dermatitis: design, validation, and scoring of the Atopic Dermatitis Control Tool (ADCT). *Curr Med Res Opin*. 2020;36(3):367-376. doi:10.1080/03007995.2019.1699516

Worst Scratch Itch Numeric Rating Scale (WSI-NRS).^{26,27}

85. Finally, Defendants have funded and promoted hundreds of continuing medical education programs as a means to disseminate seemingly unbiased unbranded disease state awareness marketing materials directly to healthcare providers to increase screening and diagnoses of atopic dermatitis, asthma and other indicated health conditions and corresponding prescriptions of Dupixent. Defendants further established ADVENT, a dedicated global medical education platform disseminating “non-promotional” information to healthcare providers worldwide regarding Type 2 inflammatory diseases that Dupixent has been approved to treat or in which Dupixent is currently being investigated.²⁸

86. In addition to extensive branded and unbranded disease awareness marketing, Defendants have also leveraged influential third parties to inappropriately characterize the safety and effectiveness of Dupixent and promote its use in an unrestricted manner inconsistent with its FDA-approved indications. In this arm of the marketing scheme, Defendants partnered with patient advocacy organizations to further drive awareness of inflammatory health conditions and engaged and funded major academic societies, trade groups and leading subject matter experts who are charged with shaping clinical treatment guidelines.

87. Eight months before securing FDA approval to market Dupixent in the United States, Defendants, Regeneron Pharmaceuticals, Inc., Sanofi-Aventis U.S. LLC and Genzyme Corporation, via the “Sanofi-Genzyme and Regeneron Alliance”, funded and coordinated the convening of a “steering committee” to discuss and provide recommendations for the management

²⁶ Paller AS, Yosipovitch G, Weidinger S, et al. Development, Psychometric Validation and Responder Definition of Worst Itch Scale in Children with Severe Atopic Dermatitis. *Dermatol Ther (Heidelb)*. 2022;12(12):2839-2850. doi:10.1007/s13555-022-00804-z

²⁷ Paller AS, Siegfried E, Marron SE, et al. Development and validation of a caregiver-reported Numeric Rating Scale for measuring worst scratch/itch in patients aged 6 months to younger than 6 years with atopic dermatitis. *J Am Acad Dermatol*. 2024;90(2):382-385. doi:10.1016/j.jaad.2023.08.104

²⁸ ADVENT, *Welcome to ADVENT!*, <https://www.adventprogram.com/us/> (last visited Dec. 9, 2025).

of atopic dermatitis, later formalizing their opinions in a September 2017 publication termed a “Multidisciplinary Consensus” in *The Journal of Allergy and Clinical Immunology: In Practice*.²⁹ Oddly, the stated rationale for convening this “steering committee” was to “develop consensus recommendations relating to the diagnosis and treatment of patients with moderate-to-severe AD in the era of biologic therapies”, even though at the time this “steering committee” met between July 2016 and January 2017, neither Dupixent nor any other biologic therapy had been approved for the treatment of atopic dermatitis. Although Dupixent was not an approved drug for any purpose at the time of the meeting, the “steering committee” nonetheless broadly advocated for its use in patients with atopic dermatitis, even beyond the indication that would eventually be approved by the FDA. Defendants used this publication as a vessel to position their experimental drug as the premier first-line therapy for patients with atopic dermatitis and to characterize its safety profile as being superior to existing therapies, including in terms of lymphoma risk, notwithstanding that it had not yet been used by any patients under real-world conditions. Indeed, in the discussion section of Defendants’ treatment guidelines publication, the authors state “for patients with moderate-to-severe AD, the [steering committee] recommends dupilumab as a first-line systemic treatment option in adults”. Of course, however, Dupixent has never been approved by the FDA as a first-line systemic treatment for atopic dermatitis, and instead is only indicated when patients first use and fail to respond to first-line topical treatments, rendering Dupixent a second-line, third-line or even fourth-line therapy. Nonetheless, making sure to drive the message home, the authors explicitly reiterate their recommendation for use of Dupixent as a “first-line systemic treatment” four (4) separate times in the publication. Additionally, the authors provide

²⁹ Boguniewicz M, Alexis AF, Beck LA, et al. Expert Perspectives on Management of Moderate-to-Severe Atopic Dermatitis: A Multidisciplinary Consensus Addressing Current and Emerging Therapies. *J Allergy Clin Immunol Pract.* 2017;5(6):1519-1531. doi:10.1016/j.jaip.2017.08.005

their opinion that all systemic immunosuppressive treatments commonly used for atopic dermatitis (cyclosporine, methotrexate, azathioprine and mycophenolate mofetil) are associated with an “increased risk” of lymphoma, while highlighting no such risk for Dupixent. The authors also stress the importance of ensuring patients have access to materials to educate themselves regarding atopic dermatitis therapies and that patient preference should play a major role in treatment selection.

88. Furthermore, a consultant for Defendants published an editorial in *The Journal of Allergy and Clinical Immunology: In Practice* to accompany and echo the recommendations of the “Multidisciplinary Consensus” article in which he characterized the use of Dupixent as “an example of precision medicine with minimal side effects and an avoidance of risks with antibiotics, systemic corticosteroids, and cytotoxic agents”.³⁰ Defendants’ consultant also noted Dupixent “is approved for use in adult patients with atopic dermatitis”, while failing to specify its designation as a second-line therapy by the FDA upon approval.

89. Sanofi is a corporate sponsor of the International Eczema Council and currently participates at the Director’s Council level through providing \$50,000 to \$99,999 annually in unrestricted funding to the organization. Through its funding of the International Eczema Council, Sanofi and its subsidiaries receive certain important benefits, including yearly meetings with International Eczema Council leadership comprised of “the researchers and clinicians who are global influencers on atopic dermatitis research and treatment”, active engagement in surveys and outreach from International Eczema Council leaders when considering policy decisions, participation in the International Eczema Council Industry Liaison Committee and the exclusive right of first refusal to participate in funding and shaping the International Eczema Council’s

³⁰ Busse WW. Are Biotherapeutics Revolutionizing Treatment of "Allergic" Diseases?. *J Allergy Clin Immunol Pract.* 2017;5(6):1517-1518. doi:10.1016/j.jaip.2017.08.017

special projects and fellowship opportunities. Despite its explicit designation as a second-line therapy in the FDA-approved prescribing information, the International Eczema Council conspicuously recommends Defendants' drug Dupixent as first-line therapy for patients with atopic dermatitis.^{31,32,33}

90. Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC are corporate sponsors of the American Academy of Dermatology Association and currently participate at the Diamond level through providing \$500,000 or more annually in funding. In exchange for funding and participating in this organization, Defendants receive access to exclusive meetings and the ability to maintain direct contact with researchers and clinicians who are global influencers on the treatment of atopic dermatitis. Despite its explicit designation as a second-line therapy in the FDA-approved prescribing information, the current American Academy of Dermatology guidelines for the management of atopic dermatitis strongly recommend Defendants' drug Dupixent as first-line therapy for atopic dermatitis.³⁴

91. Defendants similarly sponsor, fund and/or partner with multiple influential organizations that shape guidelines for the prescription of medications in the treatment of other health conditions for which Dupixent has received FDA approval, including the American College of Allergy, Asthma and Immunology, the American Academy of Allergy, Asthma and Immunology, the Allergy & Asthma Network, the American Lung Association, and the American

³¹ Simpson EL, Bruin-Weller M, Flohr C, et al. When does atopic dermatitis warrant systemic therapy? Recommendations from an expert panel of the International Eczema Council. *J Am Acad Dermatol*. 2017;77(4):623-633. doi:10.1016/j.jaad.2017.06.042

³² Drucker AM, Eyerich K, de Bruin-Weller MS, et al. Use of systemic corticosteroids for atopic dermatitis: International Eczema Council consensus statement. *Br J Dermatol*. 2018;178(3):768-775. doi:10.1111/bjd.15928

³³ Drucker AM, Lam M, Flohr C, et al. Systemic Therapy for Atopic Dermatitis in Older Adults and Adults With Comorbidities: A Scoping Review and International Eczema Council Survey. *Dermatitis*. 2022;33(3):200-206. doi:10.1097/DER.0000000000000845

³⁴ Davis DMR, Drucker AM, Alikhan A, et al. Guidelines of care for the management of atopic dermatitis in adults with phototherapy and systemic therapies. *J Am Acad Dermatol*. 2024;90(2):e43-e56. doi:10.1016/j.jaad.2023.08.102

College of Gastroenterology, among many others.

92. Defendants have also gratuitously paid Key Opinion Leaders (KOLs), distinguished researchers and clinicians with strong regional, national and international influence on treatment and prescription practices among their professional communities, to engage in marketing activities, publish research that is favorable to Dupixent and persuade fellow clinicians to prescribe Dupixent to patients with atopic dermatitis, asthma and other inflammatory health conditions.

93. According to United States Centers for Medicare & Medicaid Services (CMS) OpenPayments data, between 2019 and 2024 Defendant, Regeneron Pharmaceuticals, Inc. made 3,643 individual payments to physicians in the United States totaling \$3,041,084 for consulting fees, food and beverage costs, education costs and travel and lodging expenses associated with Dupixent. Between 2019 and 2024, Defendant's subsidiary, Regeneron Healthcare Solutions, Inc. made 420,649 individual payments to physicians in the United States totaling \$45,377,957 for consulting fees, food and beverage costs, education costs and travel and lodging expenses associated with Dupixent. Between 2019 and 2024, Defendant, Sanofi-Aventis U.S. LLC made 14,384 individual payments to physicians in the United States totaling \$3,900,992 for consulting fees, food and beverage costs, education costs and travel and lodging expenses associated with Dupixent. Between 2019 and 2024, Sanofi's subsidiary Sanofi US Services Inc. made 249 individual payments to physicians in the United States totaling \$325,906 for consulting fees, food and beverage costs, education costs and travel and lodging expenses associated with Dupixent. Finally, between 2019 and 2024, Defendant, Genzyme Corporation made 236,981 individual payments to physicians in the United States totaling \$53,277,547 for consulting fees, food and beverage costs, education costs and travel and lodging expenses associated with Dupixent.

94. Collectively, Defendants and their affiliates made 675,906 individual payments to physicians in the United States totaling \$105,923,486 for consulting fees, food and beverage costs, education costs and travel and lodging expenses associated with Dupixent between 2019 and 2024.

95. The 2017 “Multidisciplinary Consensus” publication described *supra*³⁵ provides an excellent example of how Defendants have used direct payments to KOLs to contaminate the practice of medicine for their financial gain. Ten of the 11 authors of this treatment guidelines article were retained by Defendants as physician KOLs, with the final author being a patient advocate from the National Eczema Association. In the competing interests section, Defendants disclaim any influence on the development of the treatment recommendations or the manuscript itself, and further aver “No payments were made to the authors for the writing of this manuscript”. Despite attempting to manufacture a perception that the recommendations and manuscript were free from bias, Defendants indeed financially enriched each of the authors for their work directly or indirectly associated with this manuscript. According to data from CMS OpenPayments, in 2017 Defendants paid the physician KOL authors of this manuscript a total of \$919,595 for consulting fees, food and beverage costs, education costs and travel and lodging expenses. A breakdown of these payments by author is as follows: Mark Boguniewicz, MD: \$240,959; Andrew F. Alexis, MD, MPH: \$2,102; Lisa A. Beck, MD: \$25,876; Lawrence F. Eichenfield, MD: \$100,950; Luz Fonacier, MD: \$77,589; Emma Guttman-Yassky, MD, PhD: \$91,538; Amy S. Paller, MD: \$26,556; David Pariser, MD: \$34,375; Jonathan I. Silverberg, MD, PhD, MPH: \$316,962; and Mark Lebwohl, MD: \$2,688. Furthermore, Defendants’ KOL William W. Busse, MD, author of

³⁵ Boguniewicz M, Alexis AF, Beck LA, et al. Expert Perspectives on Management of Moderate-to-Severe Atopic Dermatitis: A Multidisciplinary Consensus Addressing Current and Emerging Therapies. J Allergy Clin Immunol Pract. 2017;5(6):1519-1531. doi:10.1016/j.jaip.2017.08.005

the editorial³⁶ accompanying the “Multidisciplinary Consensus” article, was paid \$57,906 by Defendants in 2017.

96. Defendants also used KOLs to disseminate recommendations to help facilitate the transition of patients from prior effective treatments onto Dupixent, to include patient groups in which Defendants had not studied the safety and efficacy of Dupixent.³⁷

97. Further, KOLs played a pivotal role in helping Defendants push for widespread off-label use of Dupixent in the treatment of numerous unapproved inflammatory conditions including, but not limited to chronic pruritus, eczematous eruption of aging, allergic contact dermatitis, chronic hand eczema, alopecia areata, eosinophilic annular erythema and papuloerythroderma of Ofuji.³⁸

98. In addition to paying KOLs to promote and increase prescribing of Dupixent, Defendants have also disingenuously leveraged the power of their retained KOLs to sow doubt into any research shedding light on adverse effects that might lead to a reduction in prescriptions of Dupixent as further described *infra*.

99. Defendants’ aggressive tactics have paid off. In its first partial year on the market, Dupixent generated \$253.8 million in sales in the United States. In 2018, Dupixent sales in the United States climbed to \$776.3 million. Dupixent officially became a domestic blockbuster drug in 2019 based on annual sales of \$1.87 billion. United States sales of Dupixent continued to increase year over year thereafter, reaching \$3.23 billion in 2020, \$4.71 billion in 2021, \$6.67 billion in 2022, \$8.86 billion in 2023 and \$10.40 billion in 2024. To date, United States Dupixent

³⁶ Busse WW. Are Biotherapeutics Revolutionizing Treatment of "Allergic" Diseases?. J Allergy Clin Immunol Pract. 2017;5(6):1517-1518. doi:10.1016/j.jaip.2017.08.017

³⁷ Papp KA, Hong CH, Lansang MP, et al. Practical Management of Patients with Atopic Dermatitis on Dupilumab. Dermatol Ther (Heidelb). 2021;11(5):1805-1828. doi:10.1007/s13555-021-00586-w

³⁸ Hendricks AJ, Yosipovitch G, Shi VY. Dupilumab use in dermatologic conditions beyond atopic dermatitis - a systematic review. J Dermatolog Treat. 2021;32(1):19-28. doi:10.1080/09546634.2019.1689227

sales have exceeded \$30 billion.

100. Based on publicly available figures supplied by Defendants, more than an estimated 1 million patients worldwide have been treated with Dupixent across all approved indications.³⁹

101. Defendants' surreptitious marketing practices also caught the attention of the United States Department of Justice.⁴⁰ In September 2019, Defendants, Regeneron Pharmaceuticals, Inc. and Regeneron Healthcare Services, Inc. each received a civil investigative demand from the United States Department of Justice pursuant to the federal False Claims Act relating to remuneration paid to physicians in the form of consulting fees, advisory boards, speaker fees and payment or reimbursement for travel and entertainment alleged to be in violation of the federal Anti-Kickback Statute in association with several drugs including Dupixent.

E. Cutaneous T-Cell Lymphoma

102. The instant matter involves injuries of cutaneous t-cell lymphoma (CTCL) associated with the administration of Dupixent.

103. CTCL is a form of non-Hodgkin lymphoma that is initiated by the malignant proliferation of T-cell lymphocytes contained in the skin. CTCL is an extremely rare condition estimated to affect fewer than 10 out of every 1 million people in the United States annually. The disease typically presents with patches, plaques or rashes on the skin and may evolve over years.

104. CTCL is a chronic condition that is generally considered incurable, and accordingly, treatment generally aims to put the disease into remission, leaving the patient with fewer signs and symptoms. Many patients will require several different types of treatments throughout their lifetime. As lesions become more resistant to treatment, or in patients initially presenting with more severe forms of CTCL, more aggressive treatment approaches are usually

³⁹ *Atopic Dermatitis*, DUPIXENT, <https://www.dupixenthcp.com/atopicdermatitis/> (last visited Dec. 9, 2025).

⁴⁰ Annual Report 2024. Regeneron Pharmaceuticals, Inc. <https://investor.regeneron.com/pdf/2024AR>

necessary.

105. CTCL can cause severe itching, pain and physical disfigurement which can significantly impair health-related quality of life, especially in later stages of the disease. CTCL can also metastasize to affect tissues and organs other than the skin. Survival rates vary depending on stage at the time of diagnosis, and patients with moderate to advanced CTCL experience markedly reduced survival.

106. Diagnosis of CTCL is typically made through repeated skin biopsies and blood tests. Imaging and biopsies of lymph nodes and bone marrow may also be performed in certain circumstances. CTCL is graded or staged according to the TNMB (Tumor, Node, Metastasis, Blood) criteria developed by the International Society for Cutaneous Lymphomas and the European Organization for Research and Treatment of Cancer based the proportion of skin affected by lesions, the type and size of lesions, metastasis to lymph nodes or other organs and the number of Sézary cells in the blood. CTCL stages range from I to IV, and include substages indicated by letters A or B, with higher numbers and letters indicating greater severity.

107. The most common subtypes of CTCL are mycosis fungoides and Sézary syndrome. Other forms of CTCL include peripheral T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, primary cutaneous anaplastic large cell lymphoma and extranodal natural killer/T-cell lymphoma. CTCL and its subtypes fall within a broader category of malignant disease processes referred to as mature T-cell and NK-cell lymphomas.

108. CTCL is typically a non-aggressive and slowly progressing disease, taking an average of 3 to 5 years, and up to 70 years, from initial presentation of symptoms until becoming diagnosable through clinical and histopathological evaluation. Accordingly, patients who have unidentified, subclinical CTCL that goes undiagnosed or is potentially diagnosed as some other

dermatosis in the interim are likely to remain asymptomatic and live without clinical or pathological signs of CTCL for long periods of time, or even indefinitely.^{41,42,43}

109. CTCL is usually stable, and capable of remaining stable for years. However, certain environmental triggers can prompt rapid disease progression. Indeed, when undiagnosed or misdiagnosed as another dermatosis, treatment-emergent “unmasking” of CTCL has long been a known risk with the use of certain medications.⁴⁴

110. Atopic dermatitis has been a well-known and widely documented health condition, often classified under the name eczema, among the medical community since at least the 1800s.^{45,46} Although, historical accounts suggest atopic dermatitis may have been first described as early as 69-140 CE, with the first known use of the term eczema in 543 CE.⁴⁷ Early studies observed no association between atopic dermatitis and CTCL, alternatively finding that rates of atopic dermatitis were very low in CTCL patients and not significantly different from the general population.^{48,49} The sequential observation of atopic dermatitis followed by Mycosis fungoides or

⁴¹ Kim YH, Liu HL, Mraz-Gernhard S, Varghese A, Hoppe RT. Long-term outcome of 525 patients with mycosis fungoides and Sezary syndrome: clinical prognostic factors and risk for disease progression. *Arch Dermatol.* 2003;139(7):857-866. doi:10.1001/archderm.139.7.857

⁴² Lavin L, Geller S. Cutaneous T Cell Lymphoma Following Dupilumab Therapy in Patients with Atopic Dermatitis: Clinical Review and Recommendations. *Am J Clin Dermatol.* 2025;26(5):723-731. doi:10.1007/s40257-025-00955-7

⁴³ Hristov AC, Tejasvi T, Wilcox RA. Cutaneous T-cell lymphomas: 2023 update on diagnosis, risk-stratification, and management. *Am J Hematol.* 2023;98(1):193-209. doi:10.1002/ajh.26760

⁴⁴ Foo SH, Shah F, Chaganti S, Stevens A, Scarisbrick JJ. Unmasking mycosis fungoides/Sézary syndrome from preceding or co-existing benign inflammatory dermatoses requiring systemic therapies: patients frequently present with advanced disease and have an aggressive clinical course. *Br J Dermatol.* 2016;174(4):901-904. doi:10.1111/bjd.14238

⁴⁵ Crawford A. Case of Eczema Rubrum, with Remarks. *Edinb Med Surg J.* 1820;16(62):37-41.

⁴⁶ Green J. Dr. Green's Observations on the Treatment of Eczema. *Med Exam (Phila).* 1842;5(40):637-640.

⁴⁷ Kramer ON, Strom MA, Ladizinski B, Lio PA. The history of atopic dermatitis. *Clin Dermatol.* 2017;35(4):344-348. doi:10.1016/j.clindermatol.2017.03.005

⁴⁸ Tuyp E, Burgoyne A, Aitchison T, MacKie R. A case-control study of possible causative factors in mycosis fungoides. *Arch Dermatol.* 1987;123(2):196-200.

⁴⁹ Mehrany K, El-Azhary RA, Bouwhuis SA, Pittelkow MR. Cutaneous T-cell lymphoma and atopy: is there an association?. *Br J Dermatol.* 2003;149(5):1013-1017. doi:10.1111/j.1365-2133.2003.05551.x

Sézary syndrome was also extremely rare.^{50,51} However, as the use of powerful immunomodulating therapies in the treatment of dermatological conditions became commonplace, reports of CTCL in patients with atopic dermatitis – including both new-onset cases and unmasking or progression of pre-malignant cases – began to appear more frequently in the literature. Unsurprisingly, the ensuing expanded utilization of biological therapeutics in the management of dermatologic conditions was accorded by a parallel rise in CTCL diagnoses in treated patients.^{52,53,54,55} At the same time, studies reporting an increased risk of CTCL attributable to atopic dermatitis – as opposed to atopic dermatitis medications – have almost exclusively been funded and conducted by manufacturers of atopic dermatitis medications and their associates.^{56,57,58,59,60}

⁵⁰ Rajka G, Winkelmann RK. Atopic dermatitis and Sézary syndrome. *Arch Dermatol.* 1984;120(1):83-84.

⁵¹ Lange-Vejlsgaard G, Ralfkiaer E, Larsen JK, O'Connor N, Thomsen K. Fatal cutaneous T cell lymphoma in a child with atopic dermatitis. *J Am Acad Dermatol.* 1989;20(5 Pt 2):954-958. doi:10.1016/s0190-9622(89)70118-3

⁵² Sokołowska-Wojdyło M, Barańska-Rybak W, Cegielska A, Trzeciak M, Lugowska-Umer H, Gniadecki R. Atopic dermatitis-like pre-Sézary syndrome: role of immunosuppression. *Acta Derm Venereol.* 2011;91(5):574-577. doi:10.2340/00015555-1149

⁵³ Dequidt L, Franck N, Sanchez-Pena P, et al. Cutaneous lymphomas appearing during treatment with biologics: 44 cases from the French Study Group on Cutaneous Lymphomas and French Pharmacovigilance Database. *Br J Dermatol.* 2019;181(3):616-618. doi:10.1111/bjd.17834

⁵⁴ Amitay-Laish I, Guenova E, Ortiz-Romero PL, et al. The Course of Mycosis Fungoides under Cytokine Pathway Blockers: A Multicentre Analysis of Real-life Clinical Data. *Acta Derm Venereol.* 2020;100(16):adv00277. Published 2020 Sep 30. doi:10.2340/00015555-3642

⁵⁵ Davis MS, Spencer RK, Johnson CE, et al. Risk of Cutaneous T Cell Lymphoma with Psoriasis Biologic Therapies. *Dermatol Ther (Heidelb).* 2024;14(1):15-30. doi:10.1007/s13555-023-01074-z

⁵⁶ Arellano FM, Wentworth CE, Arana A, Fernández C, Paul CF. Risk of lymphoma following exposure to calcineurin inhibitors and topical steroids in patients with atopic dermatitis. *J Invest Dermatol.* 2007;127(4):808-816. doi:10.1038/sj.jid.5700622

⁵⁷ Legendre L, Barnette T, Mazereeuw-Hautier J, Meyer N, Murrell D, Paul C. Risk of lymphoma in patients with atopic dermatitis and the role of topical treatment: A systematic review and meta-analysis. *J Am Acad Dermatol.* 2015;72(6):992-1002. doi:10.1016/j.jaad.2015.02.1116

⁵⁸ Wan J, Shin DB, Syed MN, et al. Malignancy risk in patients with atopic dermatitis: a population-based cohort study. *Br J Dermatol.* 2023;189(1):53-61. doi:10.1093/bjd/ljad072

⁵⁹ Powers CM, Piontkowski AJ, Orloff J, et al. Risk of lymphoma in patients with atopic dermatitis: A case-control study in the All of Us database. *J Am Acad Dermatol.* 2024;91(2):344-346. doi:10.1016/j.jaad.2024.03.038

⁶⁰ Ma Y, Chachin M, Hirose T, et al. Prevalence and incidence of comorbidities in patients with atopic dermatitis, psoriasis, alopecia areata, and vitiligo using a Japanese claims database. *J Dermatol.* 2025;52(5):841-854. doi:10.1111/1346-8138.17643

F. Dupixent and CTCL

111. Despite its historical infrequency in the population, CTCL has been widely reported among patients taking Dupixent. These cases have included diagnoses of new, primary CTCL as well as an unexpected “unmasking” or severe and rapid clinical progression of otherwise subclinical and undiagnosed CTCL occurring weeks to years following initiation of Dupixent use. Evidence supporting a causal relationship between use of Dupixent and incident CTCL is derived from both experimental and observational studies in humans.

a. Clinical Trials

112. Years before its approval for marketing by FDA, CTCL was first reported in a patient treated with Dupixent in a Phase II clinical trial in adults with atopic dermatitis (Study R668-AD-1314). In this study, Mycosis fungoides stage IV was reported in 1 of 97 (1.03%) subjects treated with dupilumab and 0 of 97 subjects assigned to placebo.⁶¹ According to the FDA Medical Review for Dupixent, this subject presented on day 35 of treatment with apparent worsening of his atopic dermatitis after receiving 7 doses of Dupixent 300 mg. After receiving another dose on day 43, his atopic dermatitis was considered “completely refractory to treatment”, and skin biopsies were obtained. On day 48 this subject was formally diagnosed with Mycosis fungoides stage IV and Dupixent was discontinued. Mycosis fungoides stage IV was noted to be ongoing at the end of the study.⁶²

113. Following this and another CTCL clinical trial case, Defendants amended the study protocol for the ongoing Phase III open-label trial (R668-AD-1225) on January 12, 2015 to

⁶¹ Center for Drug Evaluation and Research. Medical Review: Dupixent (dupilumab). Application Number 761055. U.S. Food and Drug Administration. March 2017.

⁶² Blauvelt A, Simpson EL, Tying SK, et al. Dupilumab does not affect correlates of vaccine-induced immunity: A randomized, placebo-controlled trial in adults with moderate-to-severe atopic dermatitis. *J Am Acad Dermatol*. 2019;80(1):158-167.e1. doi:10.1016/j.jaad.2018.07.048

designate “mycosis fungoides and cutaneous T-cell dyscrasias” as adverse events of special interest (AESIs). Based on information and belief, Defendants did not likewise amend protocols for all ongoing or subsequent clinical trials in Dupixent development programs.

114. The results of the Phase III open-label extension trial were eventually posted to ClinicalTrials.gov in October 2023 and published in JAMA Dermatology in July 2024, and indicated CTCL was reported in a total of 3 patients who continued their Dupixent treatment from prior clinical trials. One additional subject treated with Dupixent developed t-cell lymphoma.⁶³ In addition to a tripling in the number of CTCL cases reported in Dupixent-treated patients in this trial since November 2014, Mycosis fungoides was also noted to be one of the 5 most common treatment-emergent adverse events leading to discontinuation in this study.⁶⁴

115. Importantly, randomized controlled trials suffer from several important limitations that restrict the generalizability of their results to real-world populations, and many patients exposed to a drug after its approval have disease features and comorbidities or use concomitant medications which would have rendered them ineligible to participate in premarket clinical trials. This limitation is particularly true for safety endpoints, as premarket clinical trials are typically too small in size and too short in duration to detect rare, serious adverse events that may be associated with the treatment. This is because premarket clinical trials are primarily designed to detect differences between groups in efficacy endpoints, with safety being a secondary consideration. The Dupixent pivotal trials were no exception.

116. Across the 3 Dupixent pivotal trials in the treatment of atopic dermatitis, 1,344 patients were randomized to Dupixent, with just 563 following the dosing schedule that was

⁶³ U.S. National Library of Medicine. NCT01949311. Open-label Study of Dupilumab in Patients With Atopic Dermatitis. <https://clinicaltrials.gov/study/NCT01949311>

⁶⁴ Beck LA, Bissonnette R, Deleuran M, et al. Dupilumab in Adults With Moderate to Severe Atopic Dermatitis: A 5-Year Open-Label Extension Study. JAMA Dermatol. 2024;160(8):805-812. doi:10.1001/jamadermatol.2024.1536

ultimately approved by the FDA. Subjects in 2 of the pivotal trials were only treated with Dupixent for up to 16 weeks, and in the third pivotal trial subjects were treated with Dupixent for up to 52 weeks. Defendants' clinical development safety database for atopic dermatitis included a total of 2,526 subjects who received at least 1 dose of Dupixent. Notably, however, while Dupixent is intended to be a chronic therapy for indefinite use, the Defendants' clinical development safety database included just 160 subjects with at least 2 years of exposure to any dose of Dupixent at the time they submitted their BLA seeking an indication in atopic dermatitis.⁶⁵

117. The safety populations in Defendants' subsequent clinical development programs for Dupixent in the treatment of other conditions were even smaller or exposed to Dupixent for shorter durations than the atopic dermatitis development program. Defendants' clinical development safety database for asthma in patients 12 years of age and older included 1,670 subjects who received at least one dose of Dupixent at the time of BLA submission. Among these, only 350 were treated with Dupixent for at least 1 year, and no subjects were treated for 2 or more years.⁶⁶ The safety database for atopic dermatitis in 12 to 17-year-olds included just 322 subjects treated with at least 1 dose of Dupixent, of which only 35 received at least 1 year of treatment and 27 received 2 years of treatment at the time of BLA submission.⁶⁷ The chronic rhinosinusitis with nasal polyps clinical development safety database included only 440 subjects treated with at least 1 dose of Dupixent at the time of BLA submission, with 83 being treated for at least 1 year and none being treated for 2 or more years.⁶⁸ Finally, Defendants' eosinophilic esophagitis safety

⁶⁵ Center for Drug Evaluation and Research. Medical Reviews: Dupixent (dupilumab). Application Number 761055. U.S. Food and Drug Administration. March 2017.

⁶⁶ Center for Drug Evaluation and Research. Approval Package: Dupixent (dupilumab). Application Number 761055Orig1s007. U.S. Food and Drug Administration. October 2018.

⁶⁷ Center for Drug Evaluation and Research. Approval Package: Dupixent (dupilumab). Application Number 761055Orig1s012. U.S. Food and Drug Administration. March 2019.

⁶⁸ Center for Drug Evaluation and Research. Approval Package: Dupixent (dupilumab). Application Number 761055Orig1s014. U.S. Food and Drug Administration. June 2019.

database included only 367 subjects who received at least 1 dose of Dupixent at the time of BLA submission. Of 182 patients who received the recommended Dupixent dosage regimen, 40 were treated for at least 1 year and none were treated for 2 or more years.⁶⁹

118. Given the small number of exposed patients and very short periods of treatment and observation in the Dupixent clinical development programs, it is not surprising that rare and serious treatment-emergent safety concerns would begin to emerge with long-term use among hundreds of thousands of patients during the postmarketing period. This phenomenon was all the more likely given that, thanks to Defendants' successful marketing scheme, the patients being treated with Dupixent in the real world were far more diverse than the populations that were enrolled in premarket clinical trials.⁷⁰

119. Indeed, following its approval for marketing in the United States in March 2017, several independent clinicians and researchers began investigating and reporting a concerning association between Dupixent use and the development of CTCL. This large and growing body of research has been published in the form of cohort studies, cross-sectional studies, analyses of postmarketing adverse events, case series and individual case reports.

b. Case Reports, Case Series and Descriptive Observational Studies

120. On November 30, 2018 a report was published in French in *Annales de Dermatologie et de Vénéréologie*.⁷¹ Published as an abstract, this report described a 49-year-old female with a history of atopic dermatitis since adolescence who experienced worsening of her disease over the preceding 6 years. Following an inadequate response to 7 months of treatment

⁶⁹ Center for Drug Evaluation and Research. Approval Package: Dupixent (dupilumab). Application Number 761055Orig1s040. U.S. Food and Drug Administration. May 2022.

⁷⁰ Papp KA, Hong CH, Lansang MP, et al. Practical Management of Patients with Atopic Dermatitis on Dupilumab. *Dermatol Ther (Heidelb)*. 2021;11(5):1805-1828. doi:10.1007/s13555-021-00586-w

⁷¹ Bozon A, Dereure O, Szablewski V, Vincent T, Du Thanh A, Gustave V. P268: Lymphome à grandes cellules ALK1 négatif CD 30+ avec atteintes ganglionnaire et cutanée sous dupilumab. *Annales de Dermatologie et de Vénéréologie*. 2018;145(12):S272-S273. doi:10.1016/j.annder.2018.09.430

with cyclosporine and 2 months of treatment with methotrexate, the patient was started on Dupixent. After 3 months of Dupixent treatment, she developed right inguinal lymph node enlargement, plaques on her neck, elbow and lower back, and a rapidly ulcerated lesion under her right nipple. Biopsy revealed CD30+ ALK1-negative cutaneous anaplastic large cell lymphoma and positron emission tomography confirmed right inguinal, cervical and right external iliac lymph node involvement. Chemotherapy was initiated to treat the patient's CTCL. In their discussion the authors noted that anaplastic large cell lymphoma was not known to be associated with atopic dermatitis, and that prior research had shown Th2 pathway involvement in oncogenesis via IL-4 and IL-13.

121. An additional case report was published on May 2, 2019, documenting CTCL in a patient treated with Dupixent in the journal *Acta Dermato-Venereologica*.⁷² The patient in this case had atopic dermatitis since early childhood, with remission in adolescence and reemergence at age 47. Due to an inadequate response to other treatments and persistent itching, Dupixent was prescribed to treat his atopic dermatitis at age 58. One month of Dupixent treatment failed to alleviate atopic dermatitis symptoms, and instead induced a flare-up of cutaneous lesions. This exacerbation of erythematous lesions prompted physicians to order a skin biopsy. Based on the results of histopathology and computed tomography scans, the patient was diagnosed with Mycosis fungoides and Dupixent was discontinued. Despite a long history of atopic dermatitis, these authors proposed that this patient may have had Mycosis fungoides that was misdiagnosed as atopic dermatitis.

122. On September 24, 2019, a case of Mycosis fungoides in a patient treated with

⁷² Chiba T, Nagai T, Osada SI, Manabe M. Diagnosis of Mycosis Fungoides Following Administration of Dupilumab for Misdiagnosed Atopic Dermatitis. *Acta Derm Venereol*. 2019;99(9):818-819. doi:10.2340/00015555-3208

Dupixent for atopic eczema was presented at the 2019 Cutaneous Lymphoma Task Force Meeting of the European Organisation for Research and Treatment of Cancer.⁷³ This subject developed widespread rash and lymphadenopathy 9 weeks after initiating Dupixent. Dupixent was discontinued, and after initial improvement, the subject experienced rapid disease progression. The subject was ultimately diagnosed with Mycosis fungoides and started on chemotherapy. Based on their assessment, the authors believed this patient may have had preexisting Mycosis fungoides which rapidly progressed upon starting treatment with Dupixent.

123. Also at the September 2019 Cutaneous Lymphoma Task Force Meeting of the European Organisation for Research and Treatment of Cancer, researchers from several universities around the world presented the results of a descriptive study investigating CTCL among patients treated with biologics; this study was formally published in the journal *Acta Dermato-Venereologica* in September 2020.⁷⁴ These researchers included all patients diagnosed with Mycosis fungoides while being treated with tumor necrosis factor-alpha inhibitors, IL-17 inhibitors, IL-12/23 inhibitors or IL-23 inhibitors and managed at Cutaneous Lymphoma Clinics, Rabin Medical Center, Israel, University of Pittsburgh, USA, University Hospital Zürich, Switzerland, Hospital Universitario 12 de Octubre, Spain, Johns Hopkins Medicine, USA and Andreas Sygros or Attikon General Hospital, Greece through June 2019. Among 19 included patients, the authors incidentally identified 2 patients who had been treated with Dupixent – in addition to other biologics meeting study inclusion criteria – for a presumed atopic dermatitis or psoriasis. These patients were retrospectively diagnosed as having preexisting Mycosis fungoides,

⁷³ Poyner E, Bacon C, Meggitt S, Weatherhead S. A case of mycosis fungoides with large cell transformation following dupilumab treatment. *Eur J Cancer*. 2019;119:S42–S43

⁷⁴ Amitay-Laish I, Guenova E, Ortiz-Romero PL, et al. The Course of Mycosis Fungoides under Cytokine Pathway Blockers: A Multicentre Analysis of Real-life Clinical Data. *Acta Derm Venereol*. 2020;100(16):adv00277. Published 2020 Sep 30. doi:10.2340/00015555-3642

and both experienced disease progression during treatment with Dupixent and other biologics. In one patient patches progressed to plaques and in the other stage III Mycosis fungoides progressed to stage IV Mycosis fungoides with blood involvement. These authors cautioned “before considering biologics for benign cutaneous inflammatory disorders, clinicians should re-think the indication, take a second look for clinical clues of MF, revise the histology or take another biopsy, and consider blood assessment, including flow cytometry”.

124. A third abstract from the September 2019 Cutaneous Lymphoma Task Force Meeting of the European Organisation for Research and Treatment of Cancer described a series of patients with newly diagnosed Mycosis fungoides after initiating biologic medications for atopic dermatitis or psoriasis.⁷⁵ One patient, a 44-year-old female who had a life-long history of atopic dermatitis, started using Dupixent in April 2018 and initially responded to the therapy. However, after 10 months of Dupixent treatment, her condition worsened, at which point a biopsy revealed Stage II mycosis fungoides. These authors advised that patients should undergo screening with biopsies before starting treatment with biologics.

125. On March 27, 2020, clinicians published a case series of 4 patients with atopic dermatitis and 3 patients with preexisting CTCL treated with Dupixent in the Journal of the American Academy of Dermatology.⁷⁶ An abstract of this study was also published in the British Journal of Dermatology in April 2020.⁷⁷ After a brief initial improvement on Dupixent, both groups of patients experienced worsening of their symptoms. The 4 atopic dermatitis patients began to develop palmoplantar desquamation, severe skin burning and pruritus, erythroderma, and

⁷⁵ Yoo J, Shah F, Velangi S, Stewart G, Hague J, Scarisbrick J. Three cases of new diagnosis of mycosis fungoides following commencement on biologic therapies for presumed psoriasis/eczema. *Eur J Cancer*. 2019;119:S41

⁷⁶ Espinosa ML, Nguyen MT, Aguirre AS, et al. Progression of cutaneous T-cell lymphoma after dupilumab: Case review of 7 patients. *J Am Acad Dermatol*. 2020;83(1):197-199. doi:10.1016/j.jaad.2020.03.050

⁷⁷ Espinosa ML, Nguyen MT, Aguirre AS, et al. P44: Dupilumab is associated with disease worsening or unmasking of cutaneous T-cell lymphoma. *Br J Dermatol*. 2020;183(4):25. doi:10.1111/bjd.19043

enlargement and thickening of plaques, and were eventually diagnosed with mycosis fungoides or CTCL (not otherwise specified) after 4 to 27 months of Dupixent treatment. Additionally, 2 of the 3 patients with preexisting CTCL being treated with Dupixent progressed from mycosis fungoides to Sézary syndrome stage IV and ultimately died due to CTCL progression. After presenting the details of these cases, these authors noted “our observations highlight the need for caution when using dupilumab in patients with atypical dermatitis presentations without previous exclusion of Cutaneous T-cell lymphoma via skin biopsy, testing for T-cell receptor gene rearrangement, and flow cytometry of the blood”, and further advised “[a]s dupilumab becomes more commonplace in the treatment of atopic dermatitis and atopic disease, we anticipate seeing a greater number of cases of unmasked Cutaneous T-cell lymphoma in patients who initially received a diagnosis of atypical atopic dermatitis”.

126. The following month in April 2020, physicians published another case report of an atopic dermatitis patient who developed CTCL while being treated with Dupixent in the *Dermatology Online Journal*.⁷⁸ Just 2 weeks after receiving his first 600 mg loading dose of Dupixent, this patient presented with an erythrodermic rash covering 95% of his body surface. Additional treatments were initiated to control the erythroderma and newly diagnosed psoriasis, but erythroderma and lymphadenopathy persisted and prompted further testing. After 3 months of Dupixent treatment, this patient was diagnosed with Sézary syndrome. The authors concluded “The temporal relationship between the initiation of dupilumab and the onset of erythroderma suggests that dupilumab was a trigger for [Sézary syndrome] in this patient” and advised “clinicians should be aware of this association for prompt diagnosis and treatment”. These authors further cautioned “Although dupilumab provides promise in the treatment of atopic and allergic

⁷⁸ Tran J, Morris L, Vu A, Duvic M. Development of Sézary syndrome following the administration of dupilumab. *Dermatol Online J.* 2020;26(4):13030/qt1m67z8sb. Published 2020 Apr 15.

conditions, clinicians should take into account its novelty and the potential for unexpected adverse events”.

127. On June 18, 2020, another case series involving Dupixent and CTCL was published in the journal *Acta Dermato-Venereologica*.⁷⁹ This report profiled the clinical courses of 1 patient with atopic dermatitis and another patient with mycosis fungoides and a history of atopic dermatitis who were both treated with Dupixent. While the patient with mycosis fungoides and atopic dermatitis experienced improvement in his symptoms after starting Dupixent, the atopic dermatitis-only patient failed to respond to Dupixent treatment over a 2-month period. This patient instead developed intense pruritus, skin plaques and palmoplantar keratosis and was ultimately diagnosed with Sézary syndrome.

128. In another case report published June 19, 2020 in *Dermatologic Therapy*, Dupixent was initiated in a patient with a 1-year history of atopic dermatitis.⁸⁰ Following 8 doses of Dupixent, the patient experienced worsening lesions, spreading plaques and the appearance of new ulcerated tumors, and was ultimately diagnosed with tumoral-stage mycosis fungoides.

129. On July 16, 2020, another case report involving CTCL in the setting of Dupixent use was published in *The Journal of Dermatology*.⁸¹ This patient had a history of atopic dermatitis since infancy which had worsened over the previous 2 years. After administering 3 doses of Dupixent, the patient presented with severe exfoliative erythroderma with superficial lymph node swelling. Following a battery of tests, the patient was diagnosed with Sézary syndrome.

130. The *Journal Cutis* published a case series on August 27, 2020 describing CTCL in

⁷⁹ Lazaridou I, Ram-Wolff C, Bouaziz JD, et al. Dupilumab Treatment in Two Patients with Cutaneous T-cell Lymphomas. *Acta Derm Venereol*. 2020;100(16):adv00271. Published 2020 Sep 30. doi:10.2340/00015555-3576

⁸⁰ Miyashiro D, Vivarelli AG, Gonçalves F, Cury-Martins J, Sanches JA. Progression of mycosis fungoides after treatment with dupilumab: A case report. *Dermatol Ther*. 2020;33(6):e13880. doi:10.1111/dth.13880

⁸¹ Umemoto N, Demitsu T, Otaki K, et al. Dupilumab therapy in Sézary syndrome misdiagnosed as atopic dermatitis: A case report. *J Dermatol*. 2020;47(10):e356-e357. doi:10.1111/1346-8138.15501

3 Dupixent-treated patients with long-standing histories of adult-onset dermatitis.⁸² After receiving 6 or fewer injections of Dupixent, each of these patients experienced an acute exacerbation of their dermatitis. Upon discontinuation of Dupixent, each patient was diagnosed with Mycosis fungoides.

131. On December 16, 2020, another case report of CTCL developing in a patient with atopic dermatitis treated with Dupixent was published in JAAD Case Reports.⁸³ Despite initial limited improvement of his symptoms, two months after initiating Dupixent the patient presented with significant worsening of his dermatitis, weight loss and marked lymphadenopathy, and was diagnosed with Mycosis fungoides stage IV. In their discussion, the authors advocated for diligent monitoring of disease course after Dupixent initiation and prompt evaluations for CTCL in patients that exhibit an inadequate response to treatment.

132. In December 2020, a case report describing a case of CTCL involving Dupixent in a 40-year-old female with a history of atopic dermatitis since childhood was published in Dermatitis.⁸⁴ Following initial improvement, this patient exhibited a flare of her atopic dermatitis symptoms after 5 months of taking Dupixent. She continued Dupixent treatment for another 8 months, during which she experienced widespread progression of lesions to 80% of her body surface area. This patient discontinued Dupixent and was ultimately diagnosed with Mycosis fungoides stage II.

133. In January 2021, a case series of 2 patients developing CTCL shortly after starting

⁸² Hollins LC, Wirth P, Fulchiero GJ Jr, Foulke GT. Long-standing dermatitis treated with dupilumab with subsequent progression to cutaneous T-cell lymphoma. *Cutis*. 2020;106(2):E8-E11. doi:10.12788/cutis.0074

⁸³ Russomanno K, Carver DeKlotz CM. Acceleration of cutaneous T-cell lymphoma following dupilumab administration. *JAAD Case Rep*. 2020;8:83-85. Published 2020 Dec 17. doi:10.1016/j.jdc.2020.12.010

⁸⁴ Ayasse M, Nelson K, Glass F, Silverberg JI. Mycosis Fungoides Unmasked by Dupilumab Treatment in a Patient With a History of Atopic Dermatitis. *Dermatitis*. 2021;32(1S):e88-e89. doi:10.1097/DER.0000000000000679

treatment with Dupixent was published in Dermatology Online Journal.⁸⁵ Both patients were treated with Dupixent for atopic dermatitis for approximately 5 months. After failing to respond to treatment, biopsies were ordered revealing Mycosis fungoides. In discussing the cases, these authors found their report to “highlight[ed] the importance of considering cutaneous lymphoma when treating inflammatory dermatoses that are recalcitrant to targeted immunotherapies”.

134. On April 22, 2021, a German clinician published a case report involving a patient who developed Mycosis fungoides large-cell anaplastic T-cell lymphoma with involvement of the supraclavicular lymph node after initiating treatment with Dupixent.⁸⁶

135. On October 11, 2021, a case report of CTCL with a fatal outcome following initiation of Dupixent was published in JAAD Case Reports.⁸⁷ The patient experienced an improvement in skin inflammatory lesions after 5 months of Dupixent treatment, but again presented to her physician 3 weeks after developing new painful, erythematous and ulcerated plaques on her breast. Testing revealed features suggestive of an anaplastic large-cell lymphoma with cutaneous and systemic involvement and Dupixent was immediately discontinued. During the following month, the patient rapidly developed a similar skin tumor on her left forearm and rapidly progressive inguinal lymph nodes. Despite achieving a partial response after 2 lines of systemic chemotherapies, the patient ultimately died of pulmonary cytomegalovirus infection associated with severe hypoxemic infiltrative fibrosis just 12 months following the onset of the first lesion on her breast. The authors considered this unfortunate case as “add[ing] to the literature of new-onset cutaneous lymphomas associated with dupilumab therapy”.

⁸⁵ Newsom M, Hrin ML, Hamid RN, et al. Two cases of mycosis fungoides diagnosed after treatment non-response to dupilumab. *Dermatol Online J.* 2021;27(1):13030/qt1133121d. Published 2021 Jan 15.

⁸⁶ Troyanova-Slavkova S. Großzellig-anaplastisches T-Zell-Lymphom und Mycosis fungoides unter der Therapie mit Dupilumab [Anaplastic Large-Cell T-Cell Lymphoma and Mycosis fungoides under Therapy with Dupilumab]. *Aktuelle Dermatologie* 2021;47(07):331-334. doi:10.1055/a-1402-9645

⁸⁷ Du-Thanh A, Gustave V, Dereure O. Lethal anaplastic large-cell lymphoma occurring in a patient treated with dupilumab. *JAAD Case Rep.* 2021;18:4-7. Published 2021 Oct 12. doi:10.1016/j.jcdr.2021.09.020

136. In October 2021, researchers from Mayo Clinic published a study of histopathologic features in 7 atopic dermatitis patients who developed atypical lymphoid infiltrate or Mycosis fungoides while taking Dupixent in *The American Journal of Dermatopathology*.⁸⁸ All 7 patients had biopsy-proven atopic dermatitis (average of 3.4 pre-treatment biopsies), reconfirmed through secondary histopathologic review at the time of the study, and none had CTCL before starting Dupixent. These patients were between 27 and 74 years of age with an average length of Dupixent treatment of 9.8 months before developing CTCL. After starting Dupixent, all 7 patients demonstrated progressive increases in the densities of atypical lymphoid infiltrates while 6 had new atypical epidermotropic lymphocytes and papillary dermal fibrosis. One patient experienced progressive worsening of symptoms 12 months post-Dupixent initiation and received a diagnosis of Mycosis fungoides stage IV. Despite chemotherapy, this patient developed systemic symptoms and lymphadenopathy, and ultimately died of complications of CTCL 6 months after her diagnosis. Based on their findings, these authors recommended serial biopsies of patients before and during treatment with Dupixent.

137. On December 27, 2021, a case report of CTCL in a 38-year-old patient with atopic dermatitis after starting Dupixent was published in *Dermatologic Therapy*.⁸⁹ This patient had been treated with topical steroids for 3 years and Dupixent for 1 year when he began developing steroid-resistant disseminated papules. Upon presentation, this patient showed no physical or histopathologic signs CTCL. Seven months later the patient experienced rapid enlargement of a nodule in his inguinal region that demonstrated lymph node involvement on positron emission

⁸⁸ Sokumbi O, Shamim H, Davis MDP, Wetter DA, Newman CC, Comfere N. Evolution of Dupilumab-Associated Cutaneous Atypical Lymphoid Infiltrates. *Am J Dermatopathol*. 2021;43(10):714-720. doi:10.1097/DAD.0000000000001875

⁸⁹ Amagai M, Ozawa M, Amagai R, et al. Nodal anaplastic large cell lymphoma with lymphomatoid papulosis following treatment of initially presumed atopic dermatitis with dupilumab: A case report. *Dermatol Ther*. 2022;35(3):e15290. doi:10.1111/dth.15290

tomography and dense infiltration of CD30+ and ALK-lymphocytes on biopsy. He was ultimately diagnosed with CD30 + ALK-nodal anaplastic large cell lymphoma and treated with chemotherapy. The authors concluded that their case indicated Dupixent could accelerate CD30+ lymphoproliferative disorders.

138. The rash of cases did not stop there, with additional case reports of new CTCL development, CTCL “unmasking” and CTCL progression in patients after starting Dupixent continuing to be published in the succeeding months. Such publications included a case report of Hodgkin’s lymphoma and peripheral T-cell lymphoma in March 2022,⁹⁰ a case report of rapidly progressing and ultimately fatal CTCL in May 2022,⁹¹ an aggressive and fatal case of primary cutaneous gamma-delta T-cell lymphoma in May 2022,⁹² a case of erythrodermic Mycosis fungoides with suspected blood involvement in August 2022⁹³ and a case of “dupilumab-associated CD8+ [Mycosis fungoides]” stage II in November 2022.⁹⁴

139. In August 2022, researchers from the Mayo Clinic published a descriptive analysis of CTCL among patients treated with biological therapies at their facility in the *International Journal of Dermatology*.⁹⁵ Over the preceding 10-year period, 17 patients treated with biologics for rheumatologic, dermatologic or gastrointestinal conditions had subsequently been diagnosed

⁹⁰ Nakazaki K, Yoshida M, Masamoto Y, et al. Discordant lymphomas of classic Hodgkin lymphoma and peripheral T-cell lymphoma following dupilumab treatment for atopic dermatitis. *Int J Hematol*. 2022;116(3):446-452. doi:10.1007/s12185-022-03330-y

⁹¹ Poyner EFM, Bacon CM, Osborne W, Frew JA, Weatherhead SC. Dupilumab unmasking cutaneous T-cell lymphoma: report of a fatal case. *Clin Exp Dermatol*. 2022;47(5):974-976. doi:10.1111/ced.15079

⁹² Ahatov R, Good AJ, Joo M, Tipton S, Goodwin B, Kelly B. A rare case of aggressive cytotoxic T-cell lymphoma in a patient on dupilumab. *JAAD Case Rep*. 2022;24:112-114. Published 2022 May 10. doi:10.1016/j.jcdr.2022.04.023

⁹³ Hashimoto M, Miyagaki T, Komaki R, Takeuchi S, Kadono T. Development of Nodular Lesions after Dupilumab Therapy in Erythrodermic Mycosis Fungoides with Interleukin-13 Receptor alpha2 Expression. *Acta Derm Venereol*. 2022;102:adv00766. Published 2022 Aug 24. doi:10.2340/actadv.v102.2234

⁹⁴ Park A, Wong L, Lang A, Kraus C, Anderson N, Elsensohn A. Dupilumab-Associated Mycosis Fungoides with a CD8+ Immunophenotype. *Dermatopathology (Basel)*. 2022;9(4):385-391. Published 2022 Nov 30. doi:10.3390/dermatopathology9040045

⁹⁵ Schaefer L, Comfere NI, Sokumbi O. Development of cutaneous T-cell lymphoma following exposure to biologic therapy: a Mayo Clinic retrospective analysis. *Int J Dermatol*. 2023;62(7):e371-e374. doi:10.1111/ijd.16386

with CTCL. Among these 17 patients, most (n=9; 53%) had been treated with Dupixent for atopic dermatitis for between 1.2 and 24 months before their CTCL diagnosis. Specific diagnoses included Mycosis fungoides, Sézary syndrome and peripheral T-cell lymphoma, and these patients were variously treated with chemotherapy, topical steroids, extracorporeal photopheresis, ultraviolet therapy, monoclonal antibodies and interferons. One of these subjects had effectively managed her long-standing atopic dermatitis with methotrexate for over 10 years and only switched to Dupixent when she no longer responded to her maintenance treatment. Within 7 months of starting Dupixent she developed peripheral T-cell lymphoma stage IV and ultimately died 8 months later. These authors advised “further prospective investigations are warranted to elucidate dupilumab’s role in the development of CTCL. This is especially relevant, given its common use in the management of AD and growing clinical indications for use including chronic urticaria, bullous pemphigoid, and alopecia areata.”

140. Still more case reports of CTCL development in patients after starting Dupixent treatment continued to be published in 2023 and 2024, including a case of angioimmunoblastic T-cell lymphoma in January 2023,⁹⁶ a case of Mycosis fungoides in March 2023,⁹⁷ a case of peripheral T-cell lymphoma “presumably associated with dupilumab” in April 2023,⁹⁸ a case of Mycosis fungoides in May 2023,⁹⁹ a case of Mycosis fungoides stage III in May 2023,¹⁰⁰ a case

⁹⁶ Choo ZY, Akinyemi AA, Cibull T, Mehlis S, Waldinger JB. Angioimmunoblastic T-cell lymphoma unmasked by treatment with dupilumab. *JAAD Case Rep.* 2023;33:87-90. Published 2023 Jan 26. doi:10.1016/j.jder.2023.01.008

⁹⁷ Yan D, Ramachandran V, Weston G, Kim RH, Latkowski JA. Diagnosing mycosis fungoides after initiation of therapy with dupilumab: a case report and literature review. *Int J Dermatol.* 2023;62(9):e500-e503. doi:10.1111/ijd.16641

⁹⁸ Shimada M, Inano S, Kitano T. T-cell lymphoma associated with dupilumab. *Ann Hematol.* 2023;102(6):1601-1602. doi:10.1007/s00277-023-05237-y

⁹⁹ O'Neill L, Duffy RF, Manders SM. Diagnosis of Mycosis Fungoides after Dupilumab Treatment for Atopic Dermatitis. *Skinmed.* 2023;21(2):105-106

¹⁰⁰ Hsieh CY, Tsai TF. Rapid progression of cutaneous T-cell lymphoma in a patient with erythroderma during dupilumab treatment, following prior sequential azathioprine, baricitinib and cyclosporine treatments. *Indian J Dermatol Venereol Leprol.* Published online May 1, 2023. doi:10.25259/IJDVL_1090_2022

of rapidly progressive CD30+ Mycosis fungoides in May 2023,¹⁰¹ a case of Sézary syndrome in July 2023,¹⁰² a case of CTCL in August 2023¹⁰³ and a case of folliculocentric Mycosis fungoides in May 2024.¹⁰⁴

141. Researchers from Mayo Clinic published a systematic review of cases of CTCL reported in patients treated with biological therapies in the American Journal of Clinical Dermatology in January 2023.¹⁰⁵ The authors ultimately identified 28 studies describing 62 patients who had developed CTCL following exposure to a biologic agent. Among these CTCL cases, Dupixent was the most commonly used biologic agent (42%), despite several others being on the market for longer and with much greater overall patient exposure.

142. In March 2023, researchers from Johns Hopkins University School of Medicine presented the results of a single center retrospective study of patients with CTCL at the Annual Meeting of the American Academy of Dermatology.¹⁰⁶ Researchers identified 411 patients with CTCL, of which 6 were found to have received Dupixent for biopsy-proven atopic dermatitis prior to being diagnosed with Mycosis fungoides or Sézary syndrome. Initial diagnosis of atopic dermatitis had been made in adulthood for 5 and in childhood in 1, and 2 patients experienced worsening of their disease after starting Dupixent. The authors concluded “[d]upilumab can be associated with worsening skin disease”.

¹⁰¹ Toker M, Srivastava P, Amin B, Wu B. Did dupilumab unmask smoldering mycosis fungoides?. JAAD Case Rep. 2023;38:11-13. Published 2023 May 30. doi:10.1016/j.jcdr.2023.05.025

¹⁰² Hamp A, Hanson J, Alhatem A, Schwartz RA. Dupilumab-Associated Sezary Syndrome. Indian J Dermatol. 2023;68(4):459-462. doi:10.4103/ijd.ijd_580_22

¹⁰³ Malick H, Wilson A, Hall M, Meier M. Cutaneous T-cell Lymphoma Progression: A Potential Dupilumab Pitfall. Cureus. 2023;15(8):e42959. Published 2023 Aug 4. doi:10.7759/cureus.42959

¹⁰⁴ Jiang SY, Yeh J, Novoa R, Wang E, Chen M. Folliculocentric Mycosis Fungoides Masquerading as Angioedema and Allergic Contact Dermatitis. J Allergy Clin Immunol Pract. 2024;12(7):1905-1906. doi:10.1016/j.jaip.2024.04.017

¹⁰⁵ Schaefer L, Comfere N, Sokumbi O. Development of Cutaneous T-Cell Lymphoma Following Biologic Treatment: A Systematic Review. Am J Clin Dermatol. 2023;24(2):153-164. doi:10.1007/s40257-022-00749-1

¹⁰⁶ Pierog O, Talluru SM, Munjal A, Weiner D, Fazal M, Ly A, Rozati S. 44588: Clinical outcomes and characteristics in CTCL patients with a diagnosis of eczema: retrospective review at a single tertiary referral center. J Am Acad Dermatol. 2023;89(3):AB136.

143. In June 2023, researchers from Saint Louis University School of Medicine, Rutgers New Jersey Medical School and Arizona College of Osteopathic Medicine published a cross-sectional study in the Archives of Dermatological Research based on 25 patients – 20 cases published in the literature and 5 from their institutions – who had developed Mycosis fungoides after starting Dupixent treatment for atopic dermatitis.¹⁰⁷ The 5 institutional patients were treated with Dupixent for an average of 8.4 months prior to developing Mycosis fungoides. Mycosis fungoides was diagnosed at stage I in 2 patients and stage II in 1 patient, while 2 patients progressed to Sézary syndrome. The average length of Dupixent treatment prior to Mycosis fungoides diagnosis among the 20 literature cases was 11.2 months. The authors advocated for “[c]lose monitoring of these patients and further investigation of the relationship between dupilumab and [Mycosis fungoides]”.

144. In August 2023, clinicians from Italy published a retrospective analysis of atopic dermatitis patients treated at two Italian tertiary care centers in The Educational Journal of the British Association of Dermatologists.¹⁰⁸ Between June 2018 and June 2023, 995 patients with atopic dermatitis had been administered Dupixent, 91.8% with a classic atopic dermatitis phenotype and 8.1% with a prurigo nodularis-like phenotype. Seven patients who were unresponsive to Dupixent or exhibited atypical clinical features discontinued treatment and were further studied through skin biopsy and T-cell receptor gamma gene rearrangement analysis. Two of these patients, males aged 55 and 85 years, developed histologically confirmed CTCL after 11 and 19 months of Dupixent treatment. One was diagnosed with Mycosis fungoides stage I and

¹⁰⁷ Hamp A, Hanson J, Schwartz RA, Lambert WC, Alhatem A. Dupilumab-associated mycosis fungoides: a cross-sectional study. Arch Dermatol Res. 2023;315(9):2561-2569. doi:10.1007/s00403-023-02652-z

¹⁰⁸ Buffon S, Alberti Violetti S, Avallone G, et al. Mycosis fungoides and Sézary syndrome following dupilumab treatment: experience of two Italian tertiary care centres. Clin Exp Dermatol. 2023;48(12):1376-1378. doi:10.1093/ced/llad277

treated with ultraviolet therapy with partial response, while the other was diagnosed with progressive Sézary syndrome stage IV and ultimately died before initiating treatment.

145. Researchers from Memorial Sloan Kettering Cancer Center published a descriptive cohort study of CTCL in Dupixent-treated patients as an abstract in the journal *Blood* in November 2023.¹⁰⁹ An updated, final version of the study was later published in May 2024 in the *Journal of the European Academy of Dermatology and Venerology*.¹¹⁰ In this study, researchers identified 27 patients presenting to their center with new Mycosis fungoides after being treated with Dupixent for atopic dermatitis. In 16 patients, atopic dermatitis was confirmed through skin biopsy, while the remaining subjects were diagnosed based on clinical features. All patients had worsening symptoms following Dupixent use, and the median duration of Dupixent use prior to CTCL diagnosis was 10.2 months. Mycosis fungoides was diagnosed as stage I (n=16), stage II (n=4), stage III (n=2) or stage IV (n=5) after Dupixent treatment. Authors remarked that the relationship between Dupixent and CTCL deserves a dedicated biological study.

146. In November 2023, researchers from The Netherlands published a case series describing lymphoid reactions mimicking CTCL in 14 patients treated with Dupixent in *JAMA Dermatology*.¹¹¹ Three of these patients were retrospectively diagnosed as having preexisting CTCL, each of which experienced clinical worsening of CTCL in response to Dupixent treatment. Despite serving as KOLs for Defendants, Sanofi, Genzyme Corporation and Regeneron Pharmaceuticals, Inc., these authors found their study highlighted a “need for caution in continuing

¹⁰⁹ Stuver R, Dusza S, Epstein-Peterson ZD, Ghione P, Johnson W, Moskowitz A, Myskowski P, Pulitzer M, Horwitz SM, Geller S. Cutaneous T-Cell Lymphoma and Dupilumab Use: A Retrospective Matched Cohort Study of Clinical Characteristics and Treatment Outcomes. *Blood*. 2023;142:6184.

¹¹⁰ Stuver R, Dusza S, Epstein-Peterson ZD, et al. Cutaneous T-cell lymphoma and dupilumab use: A retrospective matched cohort study of clinical characteristics and treatment outcomes. *J Eur Acad Dermatol Venereol*. 2025;39(2):e114-e117. doi:10.1111/jdv.20141

¹¹¹ Boesjes CM, van der Gang LF, Bakker DS, et al. Dupilumab-Associated Lymphoid Reactions in Patients With Atopic Dermatitis. *JAMA Dermatol*. 2023;159(11):1240-1247. doi:10.1001/jamadermatol.2023.3849

dupilumab treatment in patients with AD with clinical worsening and newly reported symptoms (eg, burning sensation) after initial adequate response because these patients might develop an LR”, while further warning that “these benign LRs may evolve into CTCL if dupilumab treatment is not permanently stopped”. Finally, these authors recommended histopathologic evaluation including immunohistochemical staining in specific patients prior to prescribing Dupixent.

147. In an editorial commenting on the study by Boesjes and colleagues, a dermatologist from Northwestern University Feinberg Medical School noted that the observation of a patient with both CTCL and lymphoid reaction “rais[ed] concerns that [lymphoid reaction] may not be reversible in all cases, but rather an initial step toward lymphoma”.¹¹² The author further advised “dermatologists should remain vigilant in ruling out [Mycosis fungoides], particularly in atypical presentations”, and that “[i]n such cases, skin biopsy should be the first step prior to dupilumab prescription”.

148. At the March 2024 Annual Meeting of the American Academy of Dermatology, researchers from Johns Hopkins University School of Medicine conducted and presented the results of a single center retrospective study investigating clinical presentation and outcomes of patients with CTCL following the use of biologics.¹¹³ These clinicians reviewed 148 patients with Mycosis fungoides or Sézary syndrome and a documented history of atopic dermatitis at least 2 years prior to CTCL diagnosis, identifying 19 who had undergone treatment with biologics, including adalimumab, apremilast, dupilumab (Dupixent), risankizumab, upadacitinib or ustekinumab, for an average of 2.03 years prior to their CTCL diagnosis. CTCL patients treated

¹¹² Guitart J. Dupilumab, Atopic Dermatitis, and Mycosis Fungoides-New Insights on an Evolving Story. *JAMA Dermatol.* 2023;159(11):1177-1178. doi:10.1001/jamadermatol.2023.3846

¹¹³ Fleischli A, Pierog O, Rozati S. 54697: Clinical presentation and outcomes of Cutaneous T-cell Lymphoma patients following the use of biologic therapies: retrospective review at a single tertiary referral center. *J Am Acad Dermatol.* 2024;91(3):AB26.

with biologics including Dupixent demonstrated more frequent advanced-stage CTCL (III to IV) at diagnosis (52.9% vs 15.1%; $p=0.001$) and large cell transformation (26.3% vs. 9.3%; $p=0.033$) compared to CTCL patients who had not been treated with biologics. These researchers concluded their results suggested atopic dermatitis patients treated with biologics like Dupixent could increase the risk of advanced-stage and aggressive CTCL, and that “[t]here may be patients on biologics at risk for developing or unmasking CTCL”.

149. Johns Hopkins University School of Medicine researchers presented a similar study at the 5th World Congress of Cutaneous Lymphomas in April 2024, this time comparing the presentation and outcomes of CTCL between patients treated with Dupixent and other biologic therapies.¹¹⁴ Of 148 patients diagnosed with Mycosis fungoides or Sézary syndrome between January 2011 and July 2023 and having a documented history of atopic dermatitis at least 2 years prior to CTCL diagnosis, 19 patients were identified as having been previously treated with biologics, including 9 with Dupixent only, 4 with a combination of Dupixent and other biologics and 6 with biologics other than Dupixent only. These researchers found that patients treated with Dupixent were at a significantly higher risk of advanced-stage CTCL at the time of diagnosis ($p<0.001$) and patients treated with a combination of Dupixent and other biologics were at a significantly higher risk of CTCL progression following initial diagnosis ($p=0.04$). The authors advised “[a] thorough clinical work up should be considered prior to initiating treatment with biologic agents for chronic benign inflammatory disease to establish a clear diagnosis and patients should be monitored throughout their treatment course for signs of disease transformation”.

150. Researchers from Korea and Germany investigated clinical and pathological

¹¹⁴ Fleischli A, Pierog O, Yenokyan G, Bao A, Rozati S. Abstract #183: Presentation and outcomes of cutaneous T-cell lymphoma following the use of dupilumab compared to other biologic therapies. 5th World Congress of Cutaneous Lymphomas, April 11 to 13, 2024. Pasadena, California.

features in atopic dermatitis patients who exhibited an inadequate response to Dupixent treatment and published their findings in the Journal of the European Academy of Dermatology and Venerology in April 2024.¹¹⁵ Of 371 patients treated with Dupixent for severe atopic dermatitis, 46 (12.3%) were classified as inadequate responders. Thirty-five of these patients underwent additional evaluation, and of those, 19 (54.2%) were found to have developed Mycosis fungoides following treatment with Dupixent. These authors concluded that an inadequate response to Dupixent treatment may be an indicator of early Mycosis fungoides development in patients with atopic dermatitis.

151. In an observational study published as an abstract in the British Journal of Dermatology on June 28, 2024, clinicians from the United Kingdom described their experience with CTCL development under Dupixent treatment.¹¹⁶ Between 2019 and 2022 the authors identified 5 patients in their clinical practice who had developed atypical CTCL or Sézary syndrome after starting treatment with Dupixent; 2 of the patients with atypical CTCL had biopsies performed 1-2 years prior to starting Dupixent which showed no evidence of CTCL. As a result of this concerning finding, these physicians began requiring biopsies prior to starting Dupixent in any patient with atypical eczema and “introduced peripheral blood flow cytometric evaluation as part of [their] routine dupilumab examinations and 4 months after starting treatment”. No new cases of CTCL were identified in the 6-month period after initiating the new screening protocol. Despite this, the authors concluded “we cannot exclude the possibility of CTCL developing after starting dupilumab, and we are continuing to monitor these patients”, and additionally noted “We advocate

¹¹⁵ Kook H, Gwag HE, Park SY, et al. Detecting T-cell receptor clonality in patients with severe atopic dermatitis refractory to dupilumab. *J Eur Acad Dermatol Venereol.* 2024;38(10):1939-1946. doi:10.1111/jdv.20053

¹¹⁶ Grantham H, Best K, Weatherhead S. P037 Peripheral blood flow cytometry reveals heterogeneous but benign T-cell changes in patients with severe eczema undergoing dupilumab therapy. *Br J Dermatol.* 2024;191(Suppl1): i31. doi:10.1093/bjd/ljae090.064

further investigation including biopsy and flow cytometry in atypical cases prior to dupilumab initiation, as dupilumab can cause CTCL progression in later stages”.

152. Another case report involving Mycosis fungoides developing in a 76-year-old male patient following initiation of Dupixent was published in September 2024 in *Advances in Dermatology and Allergology*.¹¹⁷ This patient had a 7-year history of atopic dermatitis confirmed through biopsy and characterized by pruritus, chronic and relapsing course of the disease, family history of atopy, xerosis, nonspecific hand eczema and pruritus after sweating. During the 10th week of Dupixent treatment the patient developed erythroderma and significant facial and lower limb edema, and was admitted for management of exacerbated skin symptoms “likely attributed to an outbreak during dupilumab treatment”. Based on the results of several tests, the patient was ultimately diagnosed with Mycosis fungoides.

153. In December 2024, researchers from Tulane University School of Medicine published a descriptive analysis of patients diagnosed with CTCL at their facility between 2021 and 2023 in the *Journal of the American Academy of Dermatology*.¹¹⁸ Of 136 new CTCL cases during the study period, 18 had been treated with Dupixent prior to their CTCL diagnosis. Most patients had adult-onset atopic dermatitis (89%) for a mean duration of 7.17 years and no prior history of treatment with immunosuppressive therapies (72%). Pre-treatment atopic dermatitis diagnoses were confirmed through biopsy in half of the patients, and the mean length of Dupixent treatment before diagnosis of CTCL was 8.1 months. Following treatment with Dupixent, 33% of patients developed erythroderma and 22% developed tumors, and CTCL was classified as stage II

¹¹⁷ Cisoń H, Cisoń W, Białynicka-Birula B, Susel M, Białynicki-Birula R, Szepietowski JC. Mycosis fungoides unveiled following dupilumab treatment in a patient with a history of atopic dermatitis. Usefulness of HFUS in monitoring skin features. A review with a case report. *Postepy Dermatol Alergol*. 2025;42(1):5-12. doi:10.5114/ada.2024.143463

¹¹⁸ Accetta J, Gioe R, Falgout L, Chastain W, Bitar C, Boh E. Cutaneous T cell lymphoma arising in patients treated with dupilumab: A case series of 18 patients. *J Am Acad Dermatol*. 2025;92(4):924-926. doi:10.1016/j.jaad.2024.10.119

or higher in 61%. Researchers concluded by commenting that their “study adds 18 patients to the body of evidence linking dupilumab with CTCL”.

154. Researchers from Memorial Sloan Kettering Cancer Center published a descriptive cohort study investigating associations between biologic therapies, atopic dermatitis and CTCL in the *Journal of the American Academy of Dermatology* in February 2025.¹¹⁹ Researchers identified all cases presenting to their cutaneous lymphoma clinical with new confirmed diagnosis of CTCL following use of Dupixent, a Janus Kinase inhibitor (JAKi) or the IL-13 inhibitor tralokinumab since 2011. Among 32 total CTCL cases, all 32 (100%) had been treated with Dupixent before being diagnosed with CTCL, 2 (6.25%) of whom had been first treated with the JAKi upadacitinib and then Dupixent before being diagnosed with CTCL. All cases had been diagnosed with atopic dermatitis (n=30) or eczematous rash (n=2), with 22 of the atopic dermatitis/eczema diagnoses being confirmed by an outside dermatologist via skin biopsy. Researchers reevaluated 5 of the 22 skin biopsies and re-confirmed 4 (80%) as pathologic atopic dermatitis, while only in 1 case they changed the pre-treatment diagnosis to Mycosis fungoides. The authors concluded that the lack of any cases of CTCL following JAKi alone or tralokinumab in their institution “challenge[s] the hypothesis that severe chronic AD is the cause of CTCL in patients exposed to dupilumab”.

155. Owing to accumulating reports describing the development and progression of CTCL in patients treated with Dupixent, independent researchers began conducting systematic reviews of published case reports and case series and publishing their findings as early as 2021. These included a systematic review of 19 cases from 10 studies published in March 2021,¹²⁰ a

¹¹⁹ Liao V, Lavin L, Pulitzer MP, Stuver R, Geller S. Diagnosis of cutaneous T-cell lymphoma following exposure to biologic agents for atopic dermatitis: A retrospective cohort study from a single tertiary cancer center. *J Am Acad Dermatol.* 2025;92(6):1394-1395. doi:10.1016/j.jaad.2025.01.088

¹²⁰ Sugaya M. Is blocking IL-4 receptor alpha beneficial for patients with mycosis fungoides or Sézary syndrome?. *J Dermatol.* 2021;48(5):e225-e226. doi:10.1111/1346-8138.15834

systematic review of 23 cases from 13 studies published in December 2021,¹²¹ a systematic review of 27 cases from 12 studies published in September 2022,¹²² a systematic review of 23 cases from 11 studies published in December 2022¹²³ and a systematic review of 30 cases from 18 studies published in December 2024.¹²⁴ In one of these systematic reviews of Dupixent CTCL case reports the authors remarked, “Interestingly, progression of [Sézary syndrome] has been frequent, considering the rarity of the disease”.¹²⁵

156. Researchers from China presented a case series of 3 patients who developed CTCL after starting Dupixent in *Frontiers in Medicine* in April 2025.¹²⁶ These patients were between 39 and 60 years of age, and each was prescribed Dupixent to treat adult-onset atopic dermatitis. All 3 patients experienced new and worsening symptoms after receiving 2 to 12 doses Dupixent and were eventually diagnosed with Mycosis fungoides (2 patients) or Sézary syndrome (1 patient). In reviewing the available scientific literature, these authors found 19 published reports involving 31 patients with atopic dermatitis who had been treated with biologics and subsequently developed CTCL, including 27 (87.10%) that were treated with Dupixent. Ultimately, these researchers concluded that their “observations suggest that dupilumab may play a role in the onset and progression of CTCL”.

157. In a May 2025 systematic review of CTCL cases reported in Dupixent users,

¹²¹ Kołkowski K, Trzeciak M, Sokołowska-Wojdyło M. Safety and Danger Considerations of Novel Treatments for Atopic Dermatitis in Context of Primary Cutaneous Lymphomas. *Int J Mol Sci*. 2021;22(24):13388. Published 2021 Dec 13. doi:10.3390/ijms222413388

¹²² Park A, Wong L, Lang A, Kraus C, Anderson N, Elsensohn A. Cutaneous T-cell lymphoma following dupilumab use: a systematic review. *Int J Dermatol*. 2023;62(7):862-876. doi:10.1111/ijd.16388

¹²³ Jfri A, Smith JS, Larocca C. Diagnosis of mycosis fungoides or Sézary syndrome after dupilumab use: A systematic review. *J Am Acad Dermatol*. 2023;88(5):1164-1166. doi:10.1016/j.jaad.2022.12.001

¹²⁴ Guo S, Wang L, Bu D, Liu F. Tumors in the setting of dupilumab use: A review of the literature. *World Allergy Organ J*. 2024;18(1):101006. Published 2024 Dec 11. doi:10.1016/j.waojou.2024.101006

¹²⁵ Sugaya M. Is blocking IL-4 receptor alpha beneficial for patients with mycosis fungoides or Sézary syndrome?. *J Dermatol*. 2021;48(5):e225-e226. doi:10.1111/1346-8138.15834

¹²⁶ Li T, Wang G, Zhang C, Wang W, Li C, Wang Y. Case Report: Cutaneous T-cell lymphoma associated with biologic therapy: three cases and a literature review. *Front Med (Lausanne)*. 2025;12:1544912. Published 2025 Apr 28. doi:10.3389/fmed.2025.1544912

researchers aggregated and described the characteristics of 124 subjects profiled across 29 different case reports, case series and cross-sectional studies.¹²⁷ Subjects had a median age of 57.3 years (SD $\pm$ 15.7) at the time of lymphoproliferative disorder diagnosis, and more cases were reported in males (n=69; 55.65%) than females (n=55; 44.35%). Most subjects were diagnosed with early-stage Mycosis fungoides (n=57; 45.97%), while a considerable proportion were diagnosed with advanced-stage Mycosis fungoides or Sézary syndrome (n=41; 33.06%). Remaining diagnoses included Mycosis fungoides or Sézary syndrome of unclear stage (n=7; 5.65%), lymphoproliferative reaction (n=13; 10.48%) and other lymphoma (n=6; 4.84%). Median time from Dupixent initiation to biopsy-confirmed lymphoproliferative disorder was 5 months. Of 121 patients initially diagnosed with atopic dermatitis, only 6 were retrospectively diagnosed as having CTCL instead of atopic dermatitis. Three additional patients were initially diagnosed with Mycosis fungoides and treated with Dupixent off-label. All 9 patients initially or retrospectively diagnosed with CTCL demonstrated clinical or histopathological progression, with 6 in the advanced stage.

158. Researchers from the University of California Davis and Virginia Commonwealth University School of Medicine performed a systematic review and data synthesis to test the hypothesis that patients taking Dupixent and other immunomodulating medications are initially diagnosed at more advanced stages of CTCL.¹²⁸ Their systematic review yielded 116 patients on dupilumab from 27 studies, 30 patients on cyclosporine from 15 studies, 7 patients on upadacitinib from 6 studies and 12,512 patients not treated with these medications from 25 studies. Using

¹²⁷ Li M, Zhao W, Lai P, Xiao Y, Wang Y. Dupilumab-associated lymphoproliferative disorders: a comprehensive review on clinicohistopathologic features and underlying mechanisms. *Curr Opin Immunol.* 2025;94:102563. doi:10.1016/j.coi.2025.102563

¹²⁸ Lee C, Aalemi A, Le N, et al. Systematic review and analysis of dupilumab, cyclosporine, and upadactinib use on the stage of mycosis fungoides presentation. *JAAD Rev.* 2026;9:147-150. doi:10.1016/j.jdrv.2026.06.009

multivariable linear regression models and treating CTCL stage gradations (IA-IVB) as a continuous outcome (coded 1-8), patients treated with Dupixent and other immunomodulating medications were initially diagnosed at more advanced stages of Mycosis fungoides compared to patients not treated with these agents, adjusting for age and sex ($p<0.0001$). Specifically, Dupixent treatment was associated with an initial Mycosis fungoides diagnosis 1.28 stages higher than no Dupixent treatment, a difference which was statistically significant ($p<0.0001$). In an earlier version of this study presented as an abstract at the American Academy of Dermatology Annual Meeting in March 2025, these researchers reported that patients on Dupixent were initially diagnosed with Mycosis fungoides at approximately 2.5 stages higher compared to patients with no Dupixent use, adjusted for age and sex ($p<0.0001$).¹²⁹

159. In May 2026, Xiao and colleagues published a retrospective study describing the characteristics of 26 patients with “dupilumab-associated CTCL” at their institution in the Journal of the American Academy of Dermatology.¹³⁰ These researchers observed that patients developing CTCL during Dupixent treatment predominantly presented with head and neck involvement ($n=17$; 65.4%), far exceeding the rate of head and neck involvement in CTCL patients not treated with Dupixent (~30%). The authors further endeavored to perform a nested case-control study to compare these 26 patients with 104 control atopic dermatitis patients also treated with Dupixent, finding that new or progressive head and neck lesions were a significant predictor of Dupixent-associated CTCL ($p<0.0001$). A supplementary literature review identified the presence of new or progressive head and neck lesions after Dupixent treatment in 15 of 35 (42.9%) published cases

¹²⁹ Lee C, Le N, Fine J, Abbott B, Wu P. 62240 Systematic Review and Analysis of Dupilumab Use on the Stage of Mycosis Fungoides Presentation. *J Am Acad Dermatol*. 2025;93(3):AB307. 10.1016/j.jaad.2025.05.1212

¹³⁰ Xiao Y, Li M, Jiang Y, Sun J, Wang Y. New or exacerbated atypical head and neck lesions as sentinel clinical features of dupilumab-associated cutaneous T-cell lymphoma. *J Allergy Clin Immunol*. 2026 (Ahead-of-Press). doi:10.1016/j.jaci.2026.04.018

with detailed clinical descriptions of anatomic patterns, suggesting the head and neck predilection pattern may not be limited to the authors' cohort. Summarizing their findings, the authors stated "We propose that dupilumab may foster a localized immune environment conducive to T-cell expansion. In susceptible individuals, premalignant or indolent malignant clones, potentially misdiagnosed as refractory AD, may be driven to expand and manifest clinically with a florid, head and neck dominant phenotype. Thus, dupilumab may not only unmask CTCL but also shape its clinical presentation by accelerating progression within a specific anatomic context."

160. As postmarketing case reports, case series and descriptive observational studies evidencing a strong risk of CTCL with Dupixent use amassed, researchers expressed concern, advised caution and strongly recommended that atopic dermatitis patients undergo multiple skin biopsies, flow-cytometry and T-cell gene rearrangement studies to rule out underlying cutaneous malignancy prior to prescribing Dupixent. Furthermore, these clinicians recommended ongoing monitoring and repeat screening through serial biopsies and other means to detect changes in pathology during the course of Dupixent treatment that may be indicative of disease transformation. Additionally, researchers advised healthcare providers to maintain a high degree of clinical suspicion for CTCL when a patient shows signs of non-responsiveness or brief, initial responsiveness to Dupixent treatment followed by exacerbation of symptoms. Despite these calls from researchers year after year, Defendants provided no such recommendations to aid in the appropriate screening and monitoring of patients to reduce or eliminate their risk of Dupixent-induced CTCL.

161. On the contrary, Defendants, via their retained KOLs, actually advocated for *increased* usage of Dupixent when the FDA-approved dosing regimen proved inadequate to relieve symptoms of atopic dermatitis. Peter A. Lio, MD, an investigator, speaker, advisory board member

and consultant for Defendants, Sanofi-Aventis U.S. LLC, Regeneron Pharmaceuticals, Inc. and Genzyme Corporation, and Vivian Y. Shi, MD, an investigator and advisor for Defendants, Sanofi-Aventis U.S. LLC, Regeneron Pharmaceuticals, Inc. and Genzyme Corporation, along with their colleague Aleski Hendricks, published an article in 2019 in the American Journal of Clinical Dermatology recommending more aggressive prescribing of Dupixent, including increased dosages, increased dosing frequency and the addition of immunosuppressive agents to existing Dupixent regimens, in circumstances when patients exhibit an inadequate or incomplete response to Dupixent treatment.¹³¹ Notably, despite claiming “[t]here were no incentives or transactions financial or otherwise relevant to this manuscript”, in just 2018 and 2019, Defendants, Regeneron Pharmaceuticals, Inc., Genzyme Corporation and Sanofi-Aventis U.S. LLC paid Peter A. Lio, MD \$306,404 in general payments and Regeneron Healthcare Solutions, Inc., Genzyme Corporation and Sanofi-Aventis U.S. LLC paid Vivian Y. Shi, MD \$69,300 in general payments. The above recommendations for more aggressive prescribing of Dupixent in patients exhibiting an inadequate or incomplete response to treatment were echoed in another “expert panel” report funded by Defendants and authored by KOLs retained by Defendants which was published in 2021.¹³²

162. Furthermore, all times relevant hereto, Defendants’ branded marketing materials and branded websites for Dupixent have expressly advertised “NO INITIAL LAB TESTING OR ONGOING LAB MONITORING” is required for prescription and treatment with

¹³¹ Hendricks AJ, Lio PA, Shi VY. Management Recommendations for Dupilumab Partial and Non-durable Responders in Atopic Dermatitis. *Am J Clin Dermatol*. 2019;20(4):565-569. doi:10.1007/s40257-019-00436-8

¹³² Papp KA, Hong CH, Lansang MP, et al. Practical Management of Patients with Atopic Dermatitis on Dupilumab. *Dermatol Ther (Heidelb)*. 2021;11(5):1805-1828. doi:10.1007/s13555-021-00586-w

Dupixent.^{133,134,135,136} Atopic dermatitis clinical treatment guidelines funded by Defendants have similarly touted “[n]o laboratory monitoring is required before initiation or during treatment” with Dupixent.^{137,138}

c. Analytic Cohort Studies

163. In addition to the extensive body of published case reports, case series and descriptive observational studies, a strong causal relationship between Dupixent and CTCL is supported by several large, high-quality analytical cohort studies of patients with atopic dermatitis and asthma. No fewer than ten analytical cohort studies investigating CTCL risk in the setting of Dupixent use and reporting an adjusted measure of association between Dupixent and incident CTCL have been published to date.

164. The first analytical cohort study (the “Owji 2023 study”) was published December 26, 2022 in *The Journal of Allergy and Clinical Immunology: In Practice* by several independent researchers from Icahn School of Medicine at Mount Sinai and Still University and a KOL retained by Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi.¹³⁹ This was a retrospective study of 9,707 subjects with atopic dermatitis treated at a single academic institution over a 5-year period.

¹³³ DUPIXENT Specialist Referral, *Dupixent Partnering With a Specialist Brochure* (PDF), https://www.dupixentspecialistreferral.com/assets/pdfs/US.DUP.24.09.0134_DUPIXENT_AD-PCP-PED_FAQ_Brochure.pdf (last visited Dec. 10, 2025).

¹³⁴ DUPIXENT, *DUP AD Patient Profile* (Aug. 2022), https://www.dupixent.com/dam/jcr:777e1c3f-5c75-4ec4-a2d4-d6eb546f7a52/DUP.22.08.0105%20DUP%20AD%20Patient%20Profile_Ore_Aug%202022%20Update.pdf (last visited Dec. 10, 2025).

¹³⁵ DUPIXENT, *Atopic Dermatitis*, <https://www.dupixenthcp.com/atopicdermatitis/> (last visited Dec. 10, 2025).

¹³⁶ DUPIXENT, *Safety/Clinical Trial, Demonstrated Long-Term Safety Profile* <https://www.dupixenthcp.com/atopicdermatitis/efficacy-safety/safety-clinical-trial> (last visited Dec. 10, 2025).

¹³⁷ Simpson EL, Bruin-Weller M, Flohr C, et al. When does atopic dermatitis warrant systemic therapy? Recommendations from an expert panel of the International Eczema Council. *J Am Acad Dermatol.* 2017;77(4):623-633. doi:10.1016/j.jaad.2017.06.042

¹³⁸ Davis DMR, Drucker AM, Alikhan A, et al. Guidelines of care for the management of atopic dermatitis in adults with phototherapy and systemic therapies. *J Am Acad Dermatol.* 2024;90(2):e43-e56. doi:10.1016/j.jaad.2023.08.102

¹³⁹ Owji S, Ungar B, Dubin DP, et al. No association between dupilumab use and short-term cancer development in atopic dermatitis patients. *J Allergy Clin Immunol Pract.* 2023;11(5):1548-1551. doi:10.1016/j.jaip.2022.12.018

Researchers fit Cox proportional hazards models to compare the risk of several different cancers between atopic dermatitis patients treated with Dupixent and atopic dermatitis patients not treated with Dupixent. Researchers observed a 267% elevated risk of CTCL among Dupixent-treated subjects (HR 3.67; 95% CI 0.89-15.21) after adjustment for age and sex. A similar result was observed when further controlling for age, sex and prior immunosuppressant use, with Dupixent being associated with a 260% increased risk of CTCL (HR 3.60; 95% CI 0.88-14.82). Despite finding no statistically significant association between Dupixent use and any of the many cancers studied, these researchers concluded that “Cutaneous T-cell lymphoma (CTCL) may, however, prove to be the exception, given the higher incidence of CTCL in exposed versus unexposed patients, with a P value approaching significance (IR, 0.92 and 0.20, respectively; adjusted HR, 3.67; P = .073). These findings are echoed by recent reports that postulate that dupilumab increases the risk for progression of new and established CTCL. Given the possible link between CTCL and dupilumab, additional studies evaluating this association are warranted.”

165. Another analytical cohort study (the “Hasan 2024 study”) was published April 6, 2024 in the Journal of the American Academy of Dermatology. In this study, researchers from the West Virginia University School of Medicine leveraged data from TriNetX, a large deidentified electronic health record (EHR) database containing clinical observations and other information on over 275 million patients, to compare the risk of CTCL and other malignancies among atopic dermatitis patients treated or not treated with Dupixent.¹⁴⁰ Researchers developed two separate models using propensity score matching to control for potential confounders. In the first model adjusting for age only, Dupixent-treated subjects exhibited over a 300% increased risk of

¹⁴⁰ Hasan I, Parsons L, Duran S, Zinn Z. Dupilumab therapy for atopic dermatitis is associated with increased risk of cutaneous T cell lymphoma: A retrospective cohort study. J Am Acad Dermatol. 2024;91(2):255-258. doi:10.1016/j.jaad.2024.03.039

developing CTCL (OR 4.10; 95% CI 2.06-8.19) compared to untreated subjects. The observed statistically significant association persisted after excluding subjects with prior disease-modifying antirheumatic drug use and controlling for age, sex, ethnicity and race, with Dupixent use increasing the risk of CTCL among atopic dermatitis patients by 220% (OR 3.20; 95% CI 1.57-6.51).

166. Also utilizing data from TriNetX, another analytical cohort study investigating CTCL risk with use of Dupixent was published August 14, 2024 in the journal *Dermatologic Therapy* by researchers from Thomas Jefferson University.¹⁴¹ In this study (the “Mandel 2024 study”), authors first excluded all subjects with a history of inflammatory conditions and use of biologics with evidence of a possible association with lymphoma. After controlling for age at the time of atopic dermatitis diagnosis, race and sex through propensity score matching, Dupixent-treated atopic dermatitis patients were found to be at a 359% increased risk of CTCL (RR 4.59; 95% CI 2.46-8.57) compared to atopic dermatitis patients not treated with Dupixent.

167. Another analytical cohort study (the “Tsai 2024 study”) was published in *The Journal of Allergy and Clinical Immunology* August 1, 2024 by researchers from Harvard Medical School, Massachusetts General Hospital, Boston Children’s Hospital and Perelman School of Medicine at the University of Pennsylvania.¹⁴² Also using data from the TriNetX database, this study followed atopic dermatitis patients over a 2-year period to compare multiple safety endpoints between those treated with Dupixent or Janus kinase inhibitors (JAKi). These researchers excluded patients with a prior diagnosis of certain rheumatic conditions other than atopic dermatitis

¹⁴¹ Mandel J, Mehta J, Hafer R, et al. Increased Risk of Cutaneous T-Cell Lymphoma Development after Dupilumab Use for Atopic Dermatitis. *Dermatol Ther.* 2024;2024:9924306. doi:10.1155/2024/9924306

¹⁴² Tsai SY, Phipatanakul W, Hawryluk EB, Oyoshi MK, Schneider LC, Ma KS. Comparative safety of oral Janus kinase inhibitors versus dupilumab in patients with atopic dermatitis: A population-based cohort study. *J Allergy Clin Immunol.* 2024;154(5):1195-1203.e3. doi:10.1016/j.jaci.2024.07.019

(alopecia areata, rheumatoid arthritis, psoriatic arthritis, ankylosing spondylitis, nonradiographic axial spondyloarthritis, ulcerative colitis, Crohn's disease, asthma, prurigo nodularis, chronic sinusitis with nasal polyposis and eosinophilic esophagitis) from their cohort, and used propensity score matching to adjust for age, sex, race, history of hypertension, type 2 diabetes, hyperlipidemia, ischemic heart disease, stroke or chronic kidney disease, comorbid atopic disease (food allergy, allergic rhinitis and asthma) and use of other immunomodulating medications (azathioprine, cyclosporine, methotrexate, and mycophenolate mofetil). Before propensity matching, Dupixent-treated patients exhibited a reduced risk of CTCL (HR 0.81; 95% CI 0.11-6.25) relative to JAKi patients. However, after propensity matching, Dupixent was associated with a 59% increased risk of CTCL (HR 1.59; 95% CI 0.15-16.67)¹⁴³ compared to JAKi, despite lacking statistical significance. A notable strength of this study was the use of JAKi-treated subjects as an active comparator population, which reduced the potential for indication bias, channeling bias and confounding from disease severity.

168. Another analytical cohort study (the "Neubauer 2024 research letter") was also conducted using TriNext and published as a short research letter in Journal of the American Academy of Dermatology in September 2024 by independent researchers from Thomas Jefferson University, Columbia University and Weill Cornell Medicine, and two KOLs retained by Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi.¹⁴⁴ As made apparent by the authors, this research letter was submitted in response to the Hasan 2024 study which reported a statistically significant increased risk of CTCL among atopic dermatitis patients treated with Dupixent. This study used a 5-year follow-up period to compare risk of CTCL among atopic dermatitis patients

¹⁴³ This study compared JAKi patients *to* Dupixent patients. These HRs and 95% CIs are the inverse of those presented in the publication, reflecting the risk of CTCL when comparing Dupixent patients *to* JAKi patients.

¹⁴⁴ Neubauer ZJK, Brunner PM, Geskin LJ, Guttman E, Lipner SR. Decoupling the association of dupilumab with cutaneous T-cell lymphoma. J Am Acad Dermatol. 2024;91(6):1296-1298. doi:10.1016/j.jaad.2024.08.057

initially treated with Dupixent or 3 alternative therapies. This study reported a non-significantly elevated risk of CTCL with Dupixent when compared to initial treatment with oral prednisone (OR 1.834; 95% CI 0.908-3.708), a non-significantly reduced risk compared to initial treatment with cyclosporine (OR 0.833; 95% CI 0.419-1.654) and a significantly reduced risk of CTCL compared to initial treatment with methotrexate (OR 0.512; 95% CI 0.295-0.892), adjusting for age, sex and race though propensity score matching. Of note, the Dupixent group in this study was comprised exclusively of patients using Dupixent off-label as a first-line, initial therapy. Relatedly, as the exposure assignment reflected only the first therapy prescribed after atopic dermatitis diagnosis without considering or adjusting for subsequent use of different medications, it is likely many subjects in the alternative therapy groups, especially in more severe cases, actually used multiple therapies including Dupixent at some point during the study period, biasing point estimates towards the null or reversing their direction entirely.

169. Another analytical cohort study (the “Sheng-Kai Ma 2025 study”) was published in June 2025 in the *European Respiratory Journal* by researchers from Harvard Medical School, Massachusetts General Hospital, Boston Children’s Hospital and Harvard T.H. Chan School of Public Health.¹⁴⁵ These researchers used data from TriNetX to compare hematological malignancy risk among asthma patients initiating treatment with Dupixent or combination inhaled corticosteroids and long-acting beta agonists (ICS/LABA) while excluding patients with a history of atopic dermatitis. Before adjustment, Dupixent-treated asthma patients had a significantly increased risk of CTCL (HR 5.97; 95% CI 2.75-12.97), Mycosis fungoides and Sézary syndrome (HR 5.62; 95% CI 2.72-11.59) and peripheral T-cell lymphoma (HR 5.61; 95% CI 2.71-11.59) compared to ICS/LABA asthma patients. A greedy nearest-neighbor propensity score matching

¹⁴⁵ Sheng-Kai Ma K, Brumbaugh B, Saff RR, et al. Dupilumab and lymphoma risk among patients with asthma: a population-based cohort study. *Eur Respir J.* 2025 (Ahead-of-Print). doi:10.1183/13993003.00139-2025

algorithm was used to control for age, sex, socioeconomic status, comorbidities, concomitant medications, substance use and healthcare utilization prior to treatment initiation. Results were nearly identical after propensity score matching, with a significant increased risk of CTCL (HR 5.63; 95% CI 1.16-27.37), Mycosis fungoides and Sézary syndrome (HR 5.79; 95% CI 1.22-27.42) and peripheral T-cell lymphoma (HR 6.14; 95% CI 1.29-29.17) among Dupixent patients. Dupixent was also associated with a higher risk of any lymphoma (HR 1.79; 95% CI 1.19-2.71) and combined mature T and NK cell lymphomas (HR 4.58; 95% CI 1.82-11.53). In a sensitivity analysis among subjects with at least 16 weeks of exposure to their respective drug, crude risk estimates comparing asthma patients treated with Dupixent to asthma patients treated with ICS/LABA increased for CTCL (HR 6.79; 95% CI 2.73-16.85) and peripheral T-cell lymphoma (HR 10.55; 95% CI 4.51-24.70) and remained similar for Mycosis fungoides and Sézary syndrome (HR 5.36; 95% CI 2.17-13.24). After propensity score matching there was an insufficient number of cases in the ICS/LABA group to calculate HRs, but log-rank tests demonstrated significantly different survival curves between dupilumab and ICS/LABA groups. The risk estimate for combined mature T and NK cell lymphomas increased considerably (HR 15.63; 95% CI 1.99-122.76) after propensity score matching in this subgroup, indicating Dupixent-treated asthma patients were at a 1,463% increased risk of these cancers compared to ICS/LABA-treated asthma patients.

170. Researchers from Columbia University, the University of Illinois, Drexel University and the University of Louisville published the results of another analytical cohort study investigating CTCL risk with Dupixent use as a research letter (the “Ahmed 2026 research letter”)

in Archives of Dermatological Research on January 3, 2026.¹⁴⁶ Using data from TriNetX, these authors created separate cohorts of patients initiating Dupixent or the JAKis upadacitinib or ruxolitinib following a diagnosis of atopic dermatitis, after excluding patients with prior diagnoses of CTCL and other malignancies. Each of the study groups were independently propensity score matched to atopic dermatitis patients not treated with a biologic medication based on age, sex, race and ethnicity, to compare CTCL risk. Significantly greater odds of incident CTCL were observed among patients treated with Dupixent compared to patients not treated with a biologic (OR 2.86; 95% CI 1.93-4.22), with 1 estimated CTCL case for every 424 Dupixent-treated patients (number needed to harm=424). In contrast, neither upadacitinib nor ruxolitinib – medications which both carry boxed warnings regarding lymphomas and other malignancies – were significantly associated with CTCL relative to non-use of biologics. Over half of the CTCL cases in Dupixent-treated patients developed >1 year after initiating treatment, suggesting unmasking of preexisting disease was unlikely to explain the association observed among this cohort. The authors concluded their findings “reinforce previous reports linking dupilumab to increased CTCL risk in AD patients”, and that “the observed association between dupilumab and CTCL, with a relatively low NNH, underscores the importance of ongoing clinical vigilance”.

171. In another analytical cohort study (the “Kridin 2026 study”) published January 20, 2026 in *Frontiers in Medicine*, the association between Dupixent and CTCL was further investigated by a KOL retained by Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC and researchers from various international facilities.¹⁴⁷ The primary aim of this

¹⁴⁶ Ahmed A, Wambold D, Burshtein J. et al. Risk of cutaneous T-cell lymphoma in atopic dermatitis patients treated with dupilumab and JAK inhibitors: a TriNetX cohort study. *Arch Dermatol Res.* 2026;318(37). doi:10.1007/s00403-025-04501-7

¹⁴⁷ Kridin K, Bieber K, Olbrich H, von Bubnoff D, Hernandez G, Zirpel H, von Bubnoff N, Thaçi D, Ludwig RJ. Dupilumab treatment is not associated with changes in lymphoma risk in atopic dermatitis and other type 2 inflammatory diseases: data from a large-scale retrospective cohort study. *Front Med.* 2026;12:1702736. doi:10.3389/fmed.2025.1702736

study was to compare lymphoma risk between patients with and without atopic dermatitis as well as patients with and without other non-dermatological type 2 inflammatory diseases including asthma, eosinophilic esophagitis and chronic sinusitis. As an apparent secondary aim, the authors also sought to assess the impact of Dupixent use on lymphoma risk among patients with atopic dermatitis and non-dermatological type 2 inflammatory diseases as compared to use of other systemic therapies. CTCL was one of seven lymphoma subtypes selected as endpoints. Although a formal study period was not specified, the authors performed multiple analyses using different exposure window definitions and controlling for different sets of variables. In maximally adjusted models, use of Dupixent for atopic dermatitis was associated with a non-significantly elevated risk of CTCL (HR 1.04; 95% CI 0.62-1.73) compared to use of other systemic atopic dermatitis therapies; the magnitude of the effect estimate increased to HR 1.41 (95% CI 0.76-2.61) when eliminating the two-month period immediately following atopic dermatitis diagnosis. Among patients with asthma, eosinophilic esophagitis or chronic sinusitis, a non-significantly reduced risk of CTCL was observed in the Dupixent group between day 1 and 4 years (HR 0.38; 95% CI 0.14-1.03) and month 3 and 4 years (HR 0.30; 95% CI 0.09-1.06) following initial non-dermatological type 2 inflammatory disease diagnosis. This study has important limitations, some of which are worth briefly highlighting. No exclusion criteria were specified (history of malignancy or otherwise), and it is unclear whether exposure status was determined by initial, any or exclusive prescription of Dupixent or other systemic therapies. Further, the index date in this study was the date atopic dermatitis or another non-dermatological type 2 inflammatory disease was diagnosed, while exposure status and outcome status were later ascertained at any time during the 4-year follow-up period. This design flaw poses at least two fatal issues: 1) It is unclear whether this study established a temporal relationship between drug exposure and the outcome because, as described,

the methods permit outcomes to occur before exposure assignment, and therefore it is possible that many lymphoma diagnoses occurred before exposure to a study medication; and 2) The study is susceptible to immortal time bias arising from person time accumulated during the interval between the index date and first drug exposure which is misattributed to the drug exposure. Because Dupixent is indicated as a second-line therapy for atopic dermatitis and an add-on therapy for asthma and chronic rhinosinusitis, it is not prescribed immediately following an initial diagnosis when used on-label. Instead, many of the medications used to define the referent groups in this study are first trialed before a prescription for Dupixent is given. As a result, person time contributed by patients in the Dupixent group during this interval has been misattributed to Dupixent exposure, diluting the observed effect of Dupixent on CTCL risk. Furthermore, abrocitinib and upadacitinib were used as exposures in the non-dermatological type 2 inflammatory disease referent group, despite the fact that neither are indicated for asthma, eosinophilic esophagitis or chronic sinusitis. Beyond the inherent imbalance in baseline risk factors caused by comparing medications indicated for use in dissimilar patient populations, as noted above, both of these medications have boxed warnings regarding risk of lymphoma and other malignancies. These factors are likely to have masked an effect of Dupixent on lymphoma risk in the non-dermatological referent group. Overall, the findings of this study would be most appropriately characterized as “mixed” or “inconclusive” given the inconsistency in the observed risks, with effect estimates falling on either side of 1.0 for CTCL and other lymphoma subtypes across analyses and most results failing to achieve significance (26 of 35 lymphoma outcomes comparing Dupixent and other systemic therapy exposure were not significant). Nonetheless, the authors mischaracterized their findings as evidence of a “decreased risk association” between Dupixent and lymphoma and as suggesting that “Treatment with dupilumab partially ameliorates

this [lymphoma] risk association”.

172. Clinicians from Orange Park Hospital in Florida, Texas A&M College of Medicine and various private dermatology clinics and medical centers in Dallas Texas published another analytical cohort study using TriNetX data as a research letter (the “Talasila 2026 research letter”) on January 23, 2026 in the International Journal of Dermatology.¹⁴⁸ These authors excluded patients with a history of immunodeficiency disorders, systemic malignancy, organ transplantation or off-study biologic exposure and used propensity scoring to match Dupixent users to non-Dupixent users based on demographic factors, immunosuppressant use and relevant comorbidities. Study cohorts were further stratified by age (18-59 years and ≥ 60 years) before comparing risk of subsequent Mycosis fungoides diagnosis by Dupixent exposure status. Among atopic dermatitis patients aged 18-59 years, Dupixent use was associated with a significant increased risk of Mycosis fungoides at 1 year (RR 2.18; 95% CI 1.08-4.40), 5 years (RR 2.38; 95% CI 1.34-4.25) and 10 years (RR 2.35; 95% CI 1.34-4.14) relative to Dupixent non-use. Dupixent users aged ≥ 60 years similarly experienced a significant increased risk of Mycosis fungoides relative to non-users at 1 year (RR 2.06; 95% CI 1.13-3.75), 5 years (RR 2.17; 95% CI 1.38-3.42) and 10 years (RR 1.97; 95% CI 1.30-2.87). Despite these findings – which were in line with prior studies on the matter – these authors chose to include certain statements that deserve further comment. First, the authors claimed “Prior studies suggest dupilumab likely unmasks or accelerates recognition of underlying CTCL, rather than inducing malignancy de novo”, and in support cited the 2023 systematic review by Park and colleagues and a 2014 atopic dermatitis management guidelines publication by

¹⁴⁸ Talasila S, Nadir U, Jeha GM, Tolkachjov SN. Diagnostic Implications of Dupilumab-Refractory Atopic Dermatitis: Risk of Underlying Cutaneous T-Cell Lymphoma and the Role of Skin Biopsy. *Int J Dermatol*. Published online January 23, 2026. doi:10.1111/ijd.70307

Eichenfield et al.¹⁴⁹ However, the Eichenfield guidelines were published over 3 years before Dupixent was even first approved for use and make no reference to Dupixent whatsoever, while Park et al. reached no such conclusion based on their literature review and instead simply acknowledged “1–2 doses of dupilumab would not be expected to have enough time to induce de novo transformation or allow enough time for unrestricted T-cell neoplastic proliferation to become clinically or histologically diagnostic of MF/CTCL”. Second, the authors concluded “The association observed in our study likely reflects diagnostic confounding and delayed recognition rather than causal carcinogenesis”. Importantly, the authors cite no support for this conclusory statement, and indeed, their data source does not permit them to make this type of inference. Instead, their findings serve to belie this theory, particularly among the 18-59 age group. Here, the relative risk of Mycosis fungoides was greater in Dupixent users at 5 years and 10 years than at 1 year, where “diagnostic confounding” and “unmasking” could theoretically be at play. Additionally, a greater proportion of Mycosis fungoides cases in the non-Dupixent group were diagnosed in year 1 than in the Dupixent group (64% v 60%), and the percentage change in the number of new cases from year 1 to year 5 (Dupixent: +58%; no Dupixent: +45%) and year 5 to year 10 (Dupixent: +5%; no Dupixent: +6%) is highly similar between groups. If “diagnostic confounding” were truly responsible for the Mycosis fungoides cases in the Dupixent group, these rates would be expected to diverge over time, with the rate of new cases declining more dramatically in the Dupixent group than the non-Dupixent group. Finally, while the author’s title calls attention to “Dupilumab-Refractory Atopic Dermatitis”, “Underlying Cutaneous T-Cell Lymphoma” and the “Role of Skin Biopsy”, none of these elements were measured, analyzed or

¹⁴⁹ Eichenfield, LF, Tom WL, Chamlin SL, Feldman SR, et al. Guidelines of care for the management of atopic dermatitis: section 1. Diagnosis and assessment of atopic dermatitis. J Am Acad Dermatol. 2014;70(2):338-351. doi:10.1016/j.jaad.2013.10.010

otherwise incorporated into this study.

173. On February 23, 2026, researchers from the University of California San Diego published the results of a retrospective single-center cohort study examining CTCL risk in relation to Dupixent use as a brief report in JAAD International (the “Baghdasarian 2026 brief report”).¹⁵⁰ Electronic health records maintained in the University of California San Diego Health system were used to identify patients with (n=1,374) or without (n=1,249,409) a history of Dupixent treatment and to ascertain incident cases of CTCL among groups, defined by diagnosis codes for Mycosis fungoides or Sézary syndrome. Patients with preexisting CTCL were excluded from the cohort and cases in the Dupixent group were not considered unless they were diagnosed at least three months after Dupixent initiation. During the study period, the absolute risk of CTCL among Dupixent and non-Dupixent patients was 0.22% and 0.042%, respectively. Using multivariable logistic regression models to adjust for age, sex, race, baseline atopic dermatitis, pain, rash, history of cutaneous squamous cell carcinoma and protocolized follow-up, these researchers found that Dupixent use was associated with 87% increased odds of CTCL (OR 1.87; 95% CI 1.08-3.23;¹⁵¹ p=0.025) compared to non-use. History of atopic dermatitis was not significantly associated with CTCL. The authors recommended vigilant baseline assessment and ongoing monitoring when initiating Dupixent, advising that physicians should exercise a low threshold for biopsy in atypical or refractory cases of atopic dermatitis and perform interval reassessments during Dupixent therapy.

174. Of the ten analytic cohort studies described above, eight reported a strong and

¹⁵⁰ Baghdasarian S, Al-Saedi O, Cuomo RE. Dupilumab and Cutaneous T-Cell Lymphoma: A Retrospective Cohort Study. JAAD Int. 2026. (Ahead-of-Press). doi:10.1016/j.jdin.2026.02.004

¹⁵¹ An odds ratio and p value were presented in the publication while a confidence interval around the odds ratio was not reported. This 95% confidence interval was constructed using data provided in Table 2 in the manuscript, operating under the presumption that “Error” reflects the standard error of the beta parameter from the logistic model.

consistent increased risk of CTCL with the use of Dupixent in patients with atopic dermatitis and asthma, with several also evidencing a positive dose-response relationship between Dupixent and CTCL. Only two of the ten cohort studies – both of which were designed, conducted and/or peer reviewed by agents retained by Defendants to perform work related to Dupixent – reported mixed findings.

175. In addition to the foregoing studies, clinicians from Johns Hopkins University School of Medicine performed a retrospective single-center cohort study to investigate the prognostic impact of prior atopic dermatitis and its treatment on CTCL and published their findings in the *British Journal of Dermatology* in April 2026.¹⁵² In the primary analysis comparing 605 CTCL patients with (n=211) and without (n=394) a prior atopic dermatitis diagnosis, history of atopic dermatitis was associated with greater odds of advanced stage CTCL at diagnosis (OR 2.05; 95% CI 1.38-3.26) and progression to advanced stage (OR 1.56; 95% CI 1.05-2.31) compared to no history of atopic dermatitis. Among the subgroup of patients with atopic dermatitis and CTCL, use of systemic therapies – as a proxy for severe atopic dermatitis – was associated with greater odds of advanced stage CTCL at diagnosis (OR 2.24; 95% CI 1.16-4.38) compared to topicals/phototherapy – as a proxy for mild to moderate atopic dermatitis. Importantly, among patients with atopic dermatitis and CTCL, Dupixent exposure was associated with even greater odds of advanced stage CTCL at diagnosis (OR 3.42; 95% CI 1.29-9.14) and progression to advanced stage CTCL (OR 3.66; 95% CI 1.40-9.80). These odds ratios reflect excess relative odds of 52.7% to 66.8% for advanced CTCL stage at diagnosis and 84.8% to 134.6% for progression to advanced stage CTCL over the relative odds observed for history of atopic dermatitis and use of

¹⁵² Khoshniyati S, Fleischli A, Niknejad K, Weiner J, Pierog O, Rozati S. The prognostic impact of prior atopic dermatitis on cutaneous T-cell lymphoma: a single-centre retrospective cohort study. *Br J Dermatol*. 2026;195(2):352-354. doi:10.1093/bjd/ljag143

any systemic therapy as a proxy for severe atopic dermatitis. In reflecting on their findings, the authors determined that misdiagnosis of early CTCL as atopic dermatitis was unlikely, commenting “Enrichment of canonical atopic comorbidities in the AD-CTCL cohort supports an atopic diathesis and reduces concern for systemic misclassification of prodromal CTCL as AD”.

d. Disproportionality Analyses

176. The FDA and all major foreign health authorities collect spontaneous reports involving adverse drug events, medication errors and product quality issues submitted by patients, healthcare providers and sponsors of pharmaceutical products. In the United States, the FDA collects and stores these reports in the FDA Adverse Event Reporting System (FAERS). It is mandatory for pharmaceutical sponsors to timely submit any adverse events it receives to the FDA. Patients and healthcare providers submit reports on a voluntary basis, either directly to the FDA or to pharmaceutical companies who are then required to forward them to the FDA. These reports are coded using the Medical Dictionary for Regulatory Activities (MedDRA), a formal, hierarchical language which assigns specific reaction preferred terms to reported adverse events.

177. Spontaneous adverse events serve as a foundation for the detection of emerging safety signals for pharmaceutical products. Pharmaceutical sponsors and independent researchers commonly analyze adverse events using statistical measures that quantify the degree of disproportionality between observed and expected reporting frequencies of drug-event combinations. Two common disproportionality statistics include the reporting odds ratio (ROR) and proportional reporting ratio (PRR), which are interpreted in a similar manner to that of the odds ratio and risk ratio, respectively. The higher the ROR or PRR, the greater the extent to which the reporting of a given adverse event is observed to be disproportionately reported with a given drug. Like odds ratios and risk ratios, RORs and PRRs are typically accompanied by 95%

confidence intervals to indicate precision and statistical significance of the disproportionality estimate. A report of an adverse event necessarily comes with it suspicion that there is an association between the drug used and event experienced. Reporters indicate the suspected role of a drug in an observed adverse event using one of the following designations: primary suspect, secondary suspect, concomitant or interacting product. A primary suspect product is one which the reporter believes is most likely responsible for the observed adverse event. A secondary suspect product is one which the reporter believes may have contributed to the observed adverse event, but which is less likely to be responsible than the drug designated the primary suspect. Concomitant products are generally not suspected to have contributed to the observed adverse event, while interacting products may have contributed to the observed adverse event by interacting with a primary or secondary suspect product.

178. To date, several published disproportionality analyses have reported a positive safety signal for CTCL in association with Dupixent based on postmarketing adverse event reports.

179. The first disproportionality analysis (the “Mota 2023 study”) was initially published as a preprint on the online preprint server Authorea on August 1, 2022 and subsequently in *Allergy: European Journal of Allergy and Clinical Immunology* on December 26, 2022 after undergoing peer review.¹⁵³ In this study, researchers from Portugal analyzed data from VigiBase, the pharmacovigilance database maintained by the World Health Organization, to investigate potential safety signals for various cancers with therapeutic biologics targeting IL-5 and IL-4 receptor alpha based on postmarketing adverse event reporting. Based on reports submitted to VigiBase between 2008 and 2020, this study identified a positive disproportionality signal for CTCL with Dupixent as reflected by a statistically significant ROR of 11.11 (95% CI 6.77-18.23).

¹⁵³ Mota D, Rama TA, Moreira A. Real-world evidence on the risk of cancer with anti-IL-5 and anti-IL-4Ra biologics. *Allergy*. 2023;78(5):1375-1377. doi:10.1111/all.15628

This finding indicates patients were over 10 times more likely to report CTCL as an adverse event with Dupixent than other medications between 2008 and 2020.

180. The second disproportionality analysis (the “Lavin 2025 study”) was published by researchers from Memorial Sloan Kettering Cancer Center in the *Journal of Investigative Dermatology* on June 28, 2024.¹⁵⁴ This study utilized data from FAERS to evaluate CTCL reporting with Dupixent and other biologics used in the treatment of dermatological conditions, and performed separate analyses of reports submitted involving a single medication (monotherapy analysis) and reports involving multiple suspect medications (polytherapy analysis). Ultimately, in analyzing adverse event reports submitted to the FDA between the first quarter of 2017 and fourth quarter of 2024, these researchers observed significant signals of disproportionate CTCL reporting with Dupixent in both monotherapy (ROR 8.81; 95% CI 7.10-10.90) and polytherapy (ROR 10.35; 95% CI 8.50-12.60) analyses. These results demonstrate Dupixent was approximately 8 to 10 times as likely to be identified as a primary suspect product in CTCL adverse event reports submitted between 2017 and 2024 as compared to other medications.

181. The third disproportionality analysis (the “Cabrera-Perez 2025 study”) also used data from FAERS and was published in *The Journal of Allergy and Clinical Immunology* by researchers from Brigham & Women’s Hospital, Harvard Medical School, Dana-Farber Cancer Institute, UMass Chan Medical School, Massachusetts General Hospital and Massachusetts Institute of Technology on November 7, 2024.¹⁵⁵ In evaluating MedDRA preferred terms for individual stages of CTCL, these researchers observed extremely strong disproportionality signals

¹⁵⁴ Lavin L, Dusza S, Geller S. Cutaneous T-Cell Lymphoma after Dupilumab Use: A Real-World Pharmacovigilance Study of the FDA Adverse Event Reporting System. *J Invest Dermatol.* 2025;145(1):211-214.e1. doi:10.1016/j.jid.2024.06.1272

¹⁵⁵ Cabrera-Perez JS, Carey VJ, Odejide OO, et al. Integrative epidemiology and immunotranscriptomics uncover a risk and potential mechanism for cutaneous lymphoma unmasking or progression with dupilumab therapy. *J Allergy Clin Immunol.* 2025;155(5):1584-1594. doi:10.1016/j.jaci.2024.10.028

with Dupixent for CTCL (PRR 29.96; 95% CI 25.02-35.87), CTCL stage I (PRR 277.00; 95% CI 118.41-648.03), CTCL stage II (PRR 103.26; 95% CI 41.20-258.82), CTCL stage III (PRR 383.54; 95% CI 115.49-1,273.73) and CTCL stage IV (PRR 1,246.52; 95% CI 281.29-5,523.87) based on reporting to FAERS between the first quarter of 2017 and fourth quarter of 2024. The findings of this disproportionality analysis indicate that various stages of CTCL were anywhere from 29 to over 1,200 times more likely to be reported to FAERS in association with Dupixent than other medications between 2017 and 2024.

182. The fourth disproportionality analysis (the “Gao 2025 study”) was published in Scientific Reports in March 2025 by researchers from Jiangyin People’s Hospital in China.¹⁵⁶ Utilizing FAERS data covering the second quarter of 2017 through the fourth quarter of 2023, this study identified significant safety signals for CTCL stage III (ROR 35.83; 95% CI 13.89-92.43) and CTCL stage IV (ROR 33.78; 95% CI 16.52-69.11) among reports in which Dupixent was identified as a primary suspect product. These results indicate Dupixent was over 30 times more likely than other medications to be identified as a primary suspect product in cases of CTCL stage III and IV reported to FAERS between 2017 and 2023.

183. In August 2025, the fifth analysis highlighting a disproportionality signal for CTCL with Dupixent (the “Zhou 2025 study”) was published in the European Journal of Pharmacology.¹⁵⁷ Researchers analyzed adverse event reporting to FAERS between the second quarter of 2017 and fourth quarter of 2024, and stratified reporting by age group (children < 18 years old and adults ≥ 18 years old). Among adults, a strong signal for CTCL stage III and CTCL

¹⁵⁶ Gao H, Cao L, Liu C. Analysis and mining of Dupilumab adverse events based on FAERS database. Sci Rep. 2025;15(1):8597. doi:10.1038/s41598-025-92330-z

¹⁵⁷ Zhou J, Xie Y, Du P, Chen M, Liu X. Analysis of differences in dupilumab-associated adverse drug event signals between children and adults based on the FAERS database. Eur J Pharmacol. 2025;1005:178103. doi:10.1016/j.ejphar.2025.178103

stage IV reporting was observed with Dupixent, as indicated by RORs of 28.10 (95% CI 13.38-59.02) and 31.11 (95% CI 14.68-65.93), respectively. Consistent with prior disproportionality analyses, the findings of this study show CTCL stage III and IV were approximately 30 times more likely to be reported to FAERS with Dupixent among adults between 2017 and 2024.

184. Additional disproportionality analyses investigating the relationship between Dupixent and CTCL continued to be published in 2026. In a 2026 study by Tanaka & Ishii, Dupixent was associated with a safety signal for cutaneous lymphoma (ROR 257.64; 95% CI 164.50-404.39) based on reporting to the Japanese Adverse Drug Event Report (JADER) database.¹⁵⁸ This signal persisted in a sensitivity analysis limited to reports submitted by physicians with available data on age and sex and no concomitant suspect products (ROR 299.41; 95% CI 178.47-502.32). Lyu and colleagues similarly reported a significant safety signal for CTCL associated with Dupixent (ROR 6.10; 95% CI 1.76-21.06), even in limiting their study to FAERS reports involving patients initiating treatments for asthma.¹⁵⁹ This signal remained after applying multivariable regression to adjust for age and sex (ROR 4.71; 95% CI 1.05-19.46).

e. Spontaneous Postmarketing Adverse Events

185. The foregoing published case reports, case series, cohort studies and disproportionality analyses were not the first sources of notice to Defendants that CTCL was a serious problem with Dupixent following its approval for marketing in the United States by the FDA. Indeed, a review of the FAERS database reveals Defendants received numerous spontaneous postmarketing adverse events of CTCL developing in patients after starting treatment with

¹⁵⁸ Tanaka H, Ishii T. Cutaneous lymphomas reported following the use of dupilumab, nemolizumab, tralokinumab, or lebrikizumab: A case investigation based on JADER database and disproportionality analysis. *Intractable Rare Dis Res.* 2026 (Ahead-of-Press). doi:10.5582/irdr.2026.01017

¹⁵⁹ Lyu W, Xu R, Chen L, et al. Neoplasm Adverse Events Associated With Anti-Type 2 Biologics: An FAERS Database Pharmacovigilance Analysis Study. *Cancer Med.* 2026;15(5):e71979. doi:10.1002/cam4.71979

Dupixent.

186. On March 6, 2018, Defendants received the first postmarketing adverse event involving CTCL associated with Dupixent use. In this case, CTCL developed in a 42-year-old female after starting Dupixent for atopic dermatitis, with CTCL coded under the MedDRA preferred terms Cutaneous T-cell lymphoma and Mycosis fungoides (Manufacturer Control # US-SA-2018SA070308; FDA Case ID 14631226). This case was submitted to Defendants by a physician in the United States, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the Mycosis fungoides reaction and the secondary suspect product in the Cutaneous T-cell lymphoma reaction.

187. On May 31, 2018, Defendants received another postmarketing adverse event involving CTCL in a 49-year-old female after starting Dupixent for atopic dermatitis, coded under the MedDRA preferred terms Cutaneous lymphoma and Anaplastic large cell lymphoma T- and null-cell types (Manufacturer Control # FR-SA-2018SA152895; FDA Case ID 14996226). This case was submitted to Defendants by a physician in France, and the outcome was coded as involving hospitalization, life-threatening and other serious outcomes. Dupixent was coded the primary suspect product in each of the CTCL reactions. In assessing causality, the reporting physician considered the patient's CTCL to be related to her use of Dupixent.

188. On June 7, 2018, Defendants received another postmarketing adverse event involving T-cell lymphoma in a 58-year-old male after starting Dupixent for atopic eczema, coded under the MedDRA preferred term T-cell lymphoma (Manufacturer Control # DE-SA-2018SA154104; FDA Case ID 15017896). This case was retrieved by Defendants from the EudraVigilance database after being submitted by a physician in Germany, and the outcome was coded as involving life-threatening outcomes. Dupixent was coded the primary suspect product in

the reaction.

189. On June 20, 2018, Defendants received another postmarketing adverse event involving T-cell lymphoma in a female patient of unspecified age after starting Dupixent, coded under the MedDRA preferred term T-cell lymphoma (Manufacturer Control # US-SA-2018SA168847; FDA Case ID 15657049). This case was submitted to Defendants by a consumer in the United States, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the reaction. Defendants determined that the T-cell lymphoma in this case was related to Dupixent.

190. On June 20, 2018, Defendants received another postmarketing adverse event involving CTCL in a female patient of unspecified age after starting Dupixent, coded under the MedDRA preferred term Cutaneous lymphoma (Manufacturer Control # US-SAKK-2018SA168812AA; FDA Case ID 15662436). This case was submitted to Defendants by a healthcare professional in the United States, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the CTCL reaction. Defendants determined that the CTCL in this case was related to Dupixent.

191. On June 28, 2018, Defendants received another postmarketing adverse event involving CTCL in a 49-year-old female after starting Dupixent for atopic dermatitis, coded under the MedDRA preferred term Cutaneous lymphoma (Manufacturer Control # FR-SA-2018SA177287; FDA Case ID 15120678). This case was retrieved by Defendants from the EudraVigilance database after being submitted by a physician in France, and the outcome was coded as involving hospitalization and other serious outcomes. Dupixent was coded the primary suspect product in the CTCL reaction.

192. On July 19, 2018, a pharmacist in the United States submitted a postmarketing

adverse event directly to the FDA involving CTCL in a 60-year-old male after starting Dupixent, coded under the MedDRA preferred term Cutaneous lymphoma (FDA Case ID 15261101). An outcome code was not provided. Dupixent was coded the primary suspect product in the CTCL reaction.

193. On August 27, 2018, Defendants received another postmarketing adverse event involving CTCL in a 71-year-old male patient after starting Dupixent for atopic dermatitis, coded under the MedDRA preferred term Cutaneous T-cell lymphoma (Manufacturer Control # US-SAKK-2018SA241856AA; FDA Case ID 15361034). This case was submitted to Defendants by a physician in the United States, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the CTCL reaction.

194. On September 21, 2018, Defendants received another postmarketing adverse event involving CTCL in a female patient of unspecified age after starting Dupixent for atopic dermatitis, coded under the MedDRA preferred terms Cutaneous T-cell lymphoma, Mycosis fungoides and T-cell lymphoma (Manufacturer Control # US-SA-2018SA266899; FDA Case ID 15453098). This case was submitted to Defendants by a healthcare professional in the United States, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in each of the CTCL reactions.

195. On October 23, 2018, Defendants received another postmarketing adverse event involving CTCL in a 69-year-old male patient after starting Dupixent for atopic dermatitis, coded under the MedDRA preferred term Cutaneous T-cell lymphoma (Manufacturer Control # DE-SA-2018SA294020; FDA Case ID 15659024). This case was submitted to Defendants by a physician in Germany, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the CTCL reaction.

196. On November 16, 2018, Defendants received another postmarketing adverse event involving CTCL in a 60-year-old male patient after starting Dupixent, coded under the MedDRA preferred term Cutaneous lymphoma (Manufacturer Control # US-SA-2018SA320596; FDA Case ID 15657860). This case was submitted to Defendants by a consumer in the United States, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the CTCL reaction.

197. Ten of the foregoing 11 spontaneous postmarketing adverse event reports of CTCL developing in patients after starting Dupixent were received by Defendants before the publication of the case report by Bozon and colleagues on November 30, 2018, with 1 additional adverse event report being submitted directly to the FDA. Defendants would have expeditiously received a copy of the single report submitted directly to the FDA if they were participants in the FDA's MedWatch-to-Manufacturer program, still in effect at the time this adverse event was reported.

198. Four additional spontaneous postmarketing adverse event reports of CTCL in patients treated with Dupixent were received by Defendants between the November 30, 2018 publication of Bozon et al. and the publication of the case report by Chiba and colleagues on May 2, 2019.

199. On January 17, 2019, Defendants sent the FDA another postmarketing adverse event it had received involving CTCL in a 62-year-old male after starting Dupixent, coded under the MedDRA preferred term Mycosis fungoides (Manufacturer Control # US-SAKK-2019SA011181AA; FDA Case ID 15837889). This case was submitted to Defendants by a physician in the United States, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the CTCL reaction.

200. On March 7, 2019, Defendants received another postmarketing adverse event

involving CTCL in a 60-year-old male after starting Dupixent for atopic eczema, coded under the MedDRA preferred terms Cutaneous T-cell lymphoma stage IV and Mycosis fungoides stage IV (Manufacturer Control # 2019SA065205). According to the narrative, this patient experienced “[r]apid progression of [M]ycosis fungoides to stage 4 disease,exacerbation of T-cell lymphoma,rapid progression of ctcl with systemic spread(developed greater than 50 cutaneous tumours)”, with “[t]umours popping up daily- very aggressive with rapid progression”, and the causal relationship between Dupixent and the reported reaction was considered “probable”. The outcome in this case was coded as involving hospitalization, disability, life-threatening and other serious outcomes, and Dupixent was coded the primary suspect product in each of the CTCL reactions.

201. On March 12, 2019, Defendants received another postmarketing adverse event involving CTCL in a male patient of unspecified age after starting Dupixent for atopic dermatitis, coded under the MedDRA preferred terms Cutaneous T-cell lymphoma and Mycosis fungoides (Manufacturer Control # CA-SA-2019SA071375; FDA Case ID 16089199). This case was submitted to Defendants by a physician in Canada, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the CTCL reaction.

202. On April 22, 2019, Defendants received another postmarketing adverse event involving CTCL in a male patient of unspecified age after starting Dupixent for asthma and allergic eczema, coded under the MedDRA preferred term Cutaneous T-cell lymphoma (Manufacturer Control # US-SA-2019SA116239; FDA Case ID 16243025). This case was submitted to Defendants by a consumer in the United States, and the outcome was coded as involving other serious outcomes. Dupixent was coded the primary suspect product in the CTCL reaction.

203. After Chiba and colleagues published their case report, spontaneous postmarketing

adverse event reports of CTCL developing in patients after starting Dupixent continued to pile up. Ten additional postmarketing adverse event reports of CTCL developing in patients after starting Dupixent were received by Defendants between May 2, 2019 and September 24, 2019, the date Poyner and colleagues presented their case report, Yoo et al. presented their case report and Amitay-Laish and colleagues presented their descriptive study at the 2019 Cutaneous Lymphoma Task Force Meeting of the European Organisation for Research and Treatment of Cancer.

204. At least 17 further postmarketing adverse event reports of CTCL developing in patients after starting Dupixent were received by Defendants and forwarded to FDA between September 24, 2019 and the publication of the case series by Espinosa and colleagues on March 27, 2020.

205. Through the end of 2020, at least 23 more postmarketing adverse event reports of CTCL developing in patients after starting Dupixent were received by Defendants and forwarded to FDA.

206. In the following years, a steady stream of CTCL adverse events in patients taking Dupixent continued to flow in. Upon information and belief, over 200 such cases were reported to the Defendants by the end of 2024.

207. These adverse event numbers have been reported, although based on well-established principles, these numbers vastly underestimate the true number of CTCL events occurring with Dupixent use. Additionally, as FDAS has made clear in its Guidance to Industry, even a single well-documented post-marketing adverse event report can constitute a safety signal requiring action by the manufacturer, including a potential label change, particularly if the report involves an event that is extremely rare in the absence of drug use.

208. Importantly, the above-referenced figures exclude any CTCL adverse event reports

associated with Dupixent that were received by Defendants via solicited reporting and/or that Defendants submitted to the FDA via periodic summary reporting and/or that Defendants failed to submit to the FDA. Thus, the true number of CTCL adverse events that have been reported to Defendants in association with Dupixent, along with the dates on which Defendants first learned of any such reports, remain unknown.

209. Many of the aforementioned adverse event reports contained causal attributions to Dupixent use by the reporting physicians.

f. Mechanistic Research

210. Defendants did not conduct any experimental carcinogenicity studies of Dupixent in any animal species prior to its approval by FDA for marketing in the United States.¹⁶⁰ Upon information and belief, since receiving approval for marketing in the United States, Defendants still have not conducted any formal testing to evaluate the carcinogenic properties of Dupixent. Similarly, upon information and belief, Defendants have failed to conduct any formal testing to identify potential mechanisms through which Dupixent exposure might cause “unmasking” or severe and rapid progression of subclinical CTCL.

211. Defendants’ failure to allocate resources towards studying causal mechanisms of Dupixent-induced CTCL is despite multiple calls to conduct such testing from independent researchers (e.g. “We believe that this relationship deserves a dedicated biological study”;^{161,162} “current hypotheses still require further validation through in vivo and in vitro experiments, as

¹⁶⁰ Center for Drug Evaluation and Research. Pharmacology Reviews: Dupixent (dupilumab). Application Number 761055. U.S. Food and Drug Administration. March 2017

¹⁶¹ Stuver R, Dusza S, Epstein-Peterson ZD, Ghione P, Johnson W, Moskowitz A, Myskowski P, Pulitzer M, Horwitz SM, Geller S. Cutaneous T-Cell Lymphoma and Dupilumab Use: A Retrospective Matched Cohort Study of Clinical Characteristics and Treatment Outcomes. *Blood*. 2023;142:6184

¹⁶² Stuver R, Dusza S, Epstein-Peterson ZD, et al. Cutaneous T-cell lymphoma and dupilumab use: A retrospective matched cohort study of clinical characteristics and treatment outcomes. *J Eur Acad Dermatol Venereol*. 2025;39(2):e114-e117. doi:10.1111/jdv.20141

well as patient sequencing data”¹⁶³) and even Defendants’ own paid KOLs (“More studies should be conducted to investigate the association of CTCL with dupilumab therapy, as the currently available data is not sufficient to identify the mechanistic relationship clearly”¹⁶⁴). Defendants’ KOLs even suggested establishing a dedicated database to provide “comprehensive data on clinical features histology, and outcomes of dupilumab-associated CTCL... given the widespread use of dupilumab”, yet upon information and belief, Defendants have failed to support any such initiative.^{165,166}

212. As a result of Defendants’ failures, the exact mechanism through which Dupixent causes or contributes to the development of CTCL has not yet been discovered. Nonetheless, a causal relationship between use of Dupixent and development or “unmasking” of CTCL is biologically plausible, and is supported by various lines of clinical and non-clinical research.

213. In the presence of inaction by Defendants to perform necessary testing and research to better understand pathological processes underlying Dupixent-induced CTCL, independent researchers and clinicians have endeavored to investigate plausible mechanisms through which Dupixent can cause CTCL or lead to “unmasking” or severe and rapid progression of subclinical CTCL.

214. IL-4 and IL-13, the two primary targets of Dupixent, have been implicated in the pathogenesis of CTCL, with IL-13 and its receptors being highly expressed in the clinically

¹⁶³ Li M, Zhao W, Lai P, Xiao Y, Wang Y. Dupilumab-associated lymphoproliferative disorders: a comprehensive review on clinicopathologic features and underlying mechanisms. *Curr Opin Immunol.* 2025;94:102563. doi:10.1016/j.coi.2025.102563

¹⁶⁴ Francuzik W, Alexiou A, Worm M. Safety of dupilumab in patients with atopic dermatitis: expert opinion. *Expert Opin Drug Saf.* 2021;20(9):997-1004. doi:10.1080/14740338.2021.1939673

¹⁶⁵ Beylot-Barry M, Staumont-Salle D. Cutaneous T-Cell Lymphoma and Dupilumab Use: A Multifactorial and Complex Story. *J Invest Dermatol.* 2025;145(1):9-11. doi:10.1016/j.jid.2024.08.015

¹⁶⁶ Amatore F, Neildez M, de Masson A, et al. Cutaneous T-cell lymphomas and dupilumab for atopic dermatitis: A systematic review and expert consensus. *J Eur Acad Dermatol Venereol.* 2026 (Ahead-of-Print). doi:10.1111/jdv.70407

involved skin of CTCL patients.^{167,168} Malignant Sézary cells have also been shown to have a T helper 2 cytokine fingerprint with a predominance of IL-13 in addition to IL-4, IL-5 and IL-10.¹⁶⁹ According to one leading theory that has been proposed by researchers, Dupixent's blocking of IL-4 receptor alpha results in increases in the amount of free IL-13, while keratinocytes which would normally act as a sink to absorb excess IL-13 in atopic conditions are also blocked by Dupixent. Consequently, the unbound IL-13 is available to bind at the upregulated IL-13 receptor alpha 2 site on cells and support proliferation of neoplastic T cell clones into full-blown lymphoma.^{170,171,172,173} This is important because prior research has identified multiple mechanisms through which free IL-13 can promote growth or survival of certain tumors through direct action on the tumor or suppression of immunosurveillance.¹⁷⁴ Clinical research has also confirmed that Dupixent administration does not lead to a reduction in IL-13 levels, despite reducing IL-4 to undetectable levels.¹⁷⁵ Of note, a similar mechanism for off-target effects of free

¹⁶⁷ Geskin LJ, Viragova S, Stolz DB, Fuschiotti P. Interleukin-13 is overexpressed in cutaneous T-cell lymphoma cells and regulates their proliferation. *Blood*. 2015;125(18):2798-2805. doi:10.1182/blood-2014-07-590398

¹⁶⁸ Mazzetto R, Miceli P, Tartaglia J, Ciolfi C, Sernicola A, Alaibac M. Role of IL-4 and IL-13 in Cutaneous T Cell Lymphoma. *Life (Basel)*. 2024;14(2):245. Published 2024 Feb 9. doi:10.3390/life14020245

¹⁶⁹ Francuzik W, Alexiou A, Worm M. Safety of dupilumab in patients with atopic dermatitis: expert opinion. *Expert Opin Drug Saf*. 2021;20(9):997-1004. doi:10.1080/14740338.2021.1939673

¹⁷⁰ Cabrera-Perez JS, Carey VJ, Odejide OO, et al. Integrative epidemiology and immunotranscriptomics uncover a risk and potential mechanism for cutaneous lymphoma unmasking or progression with dupilumab therapy. *J Allergy Clin Immunol*. 2025;155(5):1584-1594. doi:10.1016/j.jaci.2024.10.028

¹⁷¹ Guglielmo A, Borghi A, Schettini N, et al. Mycosis fungoides and IL-4/13 inhibitors: what is known and unmet needs. *Expert Rev Clin Immunol*. 2025;21(6):723-729. doi:10.1080/1744666X.2025.2507332

¹⁷² Li M, Zhao W, Lai P, Xiao Y, Wang Y. Dupilumab-associated lymphoproliferative disorders: a comprehensive review on clinicohistopathologic features and underlying mechanisms. *Curr Opin Immunol*. 2025;94:102563. doi:10.1016/j.coi.2025.102563

¹⁷³ Ong PY. Is dupilumab use in atopic dermatitis associated with cutaneous T-cell lymphoma?. *J Allergy Clin Immunol*. 2025;155(5):1481-1482. doi:10.1016/j.jaci.2025.02.002

¹⁷⁴ Geskin LJ, Viragova S, Stolz DB, Fuschiotti P. Interleukin-13 is overexpressed in cutaneous T-cell lymphoma cells and regulates their proliferation. *Blood*. 2015;125(18):2798-2805. doi:10.1182/blood-2014-07-590398

¹⁷⁵ Suleiman G, Boyango I, Dawood M, Mirmovitch-Morvay O, Avitan-Hersh E. Serum IL-13 and IL-31 Levels are not Attenuated upon Dupilumab Treatment: Lessons from a Prospective Cohort. *Acta Derm Venereol*. 2026;106:0147. doi:10.2340/actadv.v106.adv-2025-0147

IL-15 in the setting of an inadequate IL-15 sink has also been described in the literature.^{176,177} Another theory posits that pre-existing chronic skin inflammation, when exposed to a Dupixent-induced immunomodulatory shift, might promote T-cell clone emergence leading to CTCL development.¹⁷⁸ Researchers have also proposed that Dupixent may act as a trigger for the progression of initially benign lymphoid infiltrates leading to the clonal expansion of T lymphocytes and their subsequent malignant transformation.¹⁷⁹ Several additional hypotheses, including those centering around altered dendritic cell composition and cytokine alteration, among others, have also been proposed. Additionally, in rodent studies, inoculation with IL-4-producing tumor cells demonstrated rejection and long-lasting antitumor immunity, suggesting blocking IL-4 signaling may have a negative impact on tumor immunity.¹⁸⁰ Impaired antitumor responses have also been observed in mice following removal of IL-4 receptor alpha to model the effects of systemic IL-4 receptor alpha inhibition.¹⁸¹

215. More recently, researchers performed single-cell sequencing on peripheral blood mononuclear cells sampled from an atopic dermatitis patient who developed CTCL with severe blood involvement after 3 months of Dupixent treatment.¹⁸² These researchers subjected the peripheral blood mononuclear cells to treatment with IL-4 and IL-13, with and without Dupixent,

¹⁷⁶ Fehniger TA, Caligiuri MA. Interleukin 15: biology and relevance to human disease. *Blood*. 2001;97(1):14-32. doi:10.1182/blood.v97.1.14

¹⁷⁷ Ma S, Caligiuri MA, Yu J. Harnessing IL-15 signaling to potentiate NK cell-mediated cancer immunotherapy. *Trends Immunol*. 2022;43(10):833-847. doi:10.1016/j.it.2022.08.004

¹⁷⁸ Beylot-Barry M, Staumont-Salle D. Cutaneous T-Cell Lymphoma and Dupilumab Use: A Multifactorial and Complex Story. *J Invest Dermatol*. 2025;145(1):9-11. doi:10.1016/j.jid.2024.08.015

¹⁷⁹ Sokumbi O, Shamim H, Davis MDP, Wetter DA, Newman CC, Comfere N. Evolution of Dupilumab-Associated Cutaneous Atypical Lymphoid Infiltrates. *Am J Dermatopathol*. 2021;43(10):714-720. doi:10.1097/DAD.0000000000001875

¹⁸⁰ Sugaya M. Is blocking IL-4 receptor alpha beneficial for patients with mycosis fungoides or Sézary syndrome?. *J Dermatol*. 2021;48(5):e225-e226. doi:10.1111/1346-8138.15834

¹⁸¹ Yu CT, Zhao X, Marchese E, Telliou S, et al. Type 2 cytokines in breast epithelial homeostasis. *J Clin Oncol*. 2026;44:e12588-e12588. doi:10.1200/JCO.2026.44.16_suppl.e12588

¹⁸² Li M, Xiao Y, Jiang Y, Wang Y. 0912: Dupilumab unmasking cutaneous $\gamma\delta$ T cell lymphoma through promoting tumorous T cell aggressiveness and microenvironment reprogramming. *J Invest Dermatol*. 2025;145(8):S158. doi:10.1016/j.jid.2025.06.928

to explore underlying mechanisms of Dupixent and its target cytokines in CTCL pathogenesis. Dupixent was observed to reduce IL-4 receptor expression and increase IL-13 receptor alpha 1 expression in corresponding cell types. Notably, researchers found that Dupixent treatment significantly increased the proliferation of tumorous T cells, and that those tumorous T cells exhibited a markedly higher gamma-delta T-cell aggressive score and upregulated exhaustion markers. Similarly, the interferon response pathway in cells, known to be associated with aggressiveness and poor prognosis in gamma-delta T-cell lymphoma, was enriched after Dupixent treatment. Furthermore, Dupixent was observed to upregulate several genes in monocytes that are known to promote tumor progression. Ultimately, these researchers determined their case served to provide “evidence that dupilumab promotes tumor progression in T cell lymphoma”.

216. Moreover, researchers have recently proposed that Dupixent exposure can more broadly lead to unmasking of pre-existing, subclinical autoimmune disease processes beyond CTCL.^{183,184} For example, commenting on the results of a cohort study investigating incident autoimmune diseases in children with atopy treated with Dupixent, a clinician from University of Córdoba in Spain noted that the results “suggest that dupilumab may differentially influence disease trajectories depending on the underlying immunologic context” and that “targeted blockade of type 2 pathways may alter the balance, timing, or clinical expression of immune-mediated conditions across individuals”.¹⁸⁵ These observations lend further support to the probability that systemic effects of Dupixent on the immune environment are capable of provoking, aggravating, exacerbating, augmenting and/or accelerating latent, subclinical diseases,

¹⁸³ Gabryszewski SJ, Hirabayashi K, Botdorf M, et al. Incidence of autoimmune diseases in children with atopy treated with dupilumab. *J Allergy Clin Immunol*. 2026. (Ahead-of-Press). doi:10.1016/j.jaci.2026.03.010

¹⁸⁴ Fu F, Zhao Q, Ma L, Wang A, Wang Z, Xu X. Case Report: Dupilumab-associated ulcerative colitis: elucidating the pathomechanistic link between Th2 blockade and Th17 polarized intestinal inflammation. *Front. Immunol*. 2026;17:1875748. doi:10.3389/fimmu.2026.1875748

¹⁸⁵ Ruano J. When blocking type 2 immunity reveals autoimmunity: Lessons from dupilumab in pediatric atopy. *J Allergy Clin Immunol*. 2026. (Ahead-of-Press). doi:10.1016/j.jaci.2026.06.002

including CTCL.

217. The aforementioned data and information that was received by Defendants prior to Plaintiff's use of Dupixent as outlined in this Complaint constitutes newly acquired information pursuant to C.F.R. § 601.12 (f)(6), which supported a label change to Dupixent in order to properly warn about CTCL with Dupixent use.

g. Expert Consensus Statements

218. Acknowledging the rapidly expanding body of research concerning Dupixent and CTCL, professional societies and expert panels began convening to review and characterize the available evidence and issue recommendations regarding appropriate screening, diagnosis and treatment measures.

219. In 2025, the French Study Group on Cutaneous Lymphoma and the French Study Group on Atopic Eczema collaborated to develop expert consensus recommendations to “limit the use of dupilumab in ‘CTCL high-risk’ patients and guide clinicians and pathologists in the diagnosis and management of CTCL arising during dupilumab therapy”.¹⁸⁶ This group performed a systematic literature review to obtain and synthesize the available evidence concerning Dupixent and CTCL and appointed a multidisciplinary panel of 37 experts in CTCL and/or atopic dermatitis to develop draft guidelines. In total, data from 51 studies encompassing 547 patients who developed CTCL during Dupixent treatment were synthesized. The expert panel reviewed the data collected in the systematic literature review and conducted two rounds of voting on the draft statements, ultimately reaching broad consensus that “dupilumab may unmask or exacerbate pre-existing CTCL and should be avoided in cases of [Mycosis fungoides/Sézary syndrome]”. The

¹⁸⁶ Amatore F, Neildez M, de Masson A, et al. Cutaneous T-cell lymphomas and dupilumab for atopic dermatitis: A systematic review and expert consensus. *J Eur Acad Dermatol Venereol*. 2026 (Ahead-of-Print). doi:10.1111/jdv.70407

expert panel further emphasized that “patients with late-onset or atypical atopic dermatitis should undergo thorough diagnostic evaluation, including histology and clonality testing, before and during dupilumab therapy”, and recommended Dupixent be immediately discontinued upon diagnosing CTCL. The group hypothesized that “dupilumab may instigate the underlying mechanism of action, while a pro-inflammatory skin environment might enhance susceptibility to lymphomagenesis or unmask latent disease”. While acknowledging prior research suggesting an increased background risk of CTCL in patients with atopic dermatitis, these experts determined “dupilumab seems to have an additional impact”. Ultimately, the expert panel concluded “this systematic review and expert consensus underscore the need for greater caution when using dupilumab in patients at risk for CTCL. Current evidence suggests that dupilumab may unmask latent disease or worsen existing CTCL and therefore should be initiated only in cases of typical AD. Clinicians should perform comprehensive diagnostic evaluations in adults presenting with atypical, treatment-resistant or late-onset dermatitis, both before initiating and during dupilumab therapy.”

220. A second set of consensus recommendations, from the European Organization for Research and Treatment of Cancer (EORTC) Cutaneous Lymphoma Tumour Group, were subsequently published.^{187,188} A “focused narrative review of the literature” was performed to develop a “scoping questionnaire comprising 14 predefined clinical statements” that was then circulated to unspecified EORTC Cutaneous Lymphoma Tumour Group members for feedback

¹⁸⁷ Calvão J, de Masson A, Cozzio A, et al. Dupilumab therapy in atopic dermatitis when cutaneous lymphoma is suspected: Consensus recommendations from the EORTC Cutaneous Lymphoma Tumour Group. *J Allergy Clin Immunol.* 2026;157(6):1237-1245. doi:10.1016/j.jaci.2026.04.002

¹⁸⁸ The EORTC Cutaneous Lymphoma Tumour Group consensus recommendations were later presented as an abstract at the 106th Annual Meeting of the British Association of Dermatologists in June-July 2026 (doi:10.1093/bjd/ljag086.557). Dr. Patrick Brunner, who was listed as a co-author of the manuscript published in the *Journal of Allergy and Clinical Immunology*, and for whom EORTC membership status could not be determined, was omitted from the abstract presented at the 106th Annual Meeting of the British Association of Dermatologists. See §G *infra* for further discussion of Dr. Patrick Brunner.

and voting. No details regarding individual author contributions, the parameters of the non-systematic literature review or the evidence synthesis methodology were provided, and the actual recommendations were only briefly discussed in the article. Instead, much of the text in this 9-page manuscript was dedicated to defending the purported utility of Dupixent in the treatment of patients with atopic dermatitis, mischaracterizing the results of epidemiologic studies and suggesting most observed CTCL cases were misdiagnosed as atopic dermatitis. Further, the article featured unusually repetitive refrains to a lack of or inability to establish causality (e.g. “a causal relationship remains unproven”, “the absence of proven causality”, “cannot establish causality”, “cannot establish causality”, “causality remains unproven”, “limiting causal inference”, “cannot establish causality”, “True drug effect (rare, unproven)”), in a manner otherwise quite unconventional to epidemiologic research. Despite these limitations and irregularities, this group did ultimately recommend “clinicians should maintain a low threshold for reassessment in atypical, adult-onset, or treatment-resistant presentations, and/or when clinical response is absent or unexpected...Where CTCL is suspected or cannot be confidently excluded, clinicopathologic correlation—including skin biopsy and additional diagnostic testing when available—is essential, and dupilumab should generally be avoided in suspected or confirmed MF or SS”.

G. Defendants’ Reaction to Negative Research

221. In response to research identifying a strong causal connection between Dupixent and CTCL, Defendants did not express concern for patient safety or commit to taking necessary actions to protect patients and develop a better understanding of this relationship. Instead, Defendants launched an aggressive publishing campaign to assuage fears among patients and healthcare providers that might otherwise lead to a reduction in prescriptions for Dupixent. Defendants’ employees, along with KOLs paid by Defendants, published original studies and

wrote multiple letters to editors of journals that had published research concerning Dupixent and CTCL in order to sow doubt into the findings of these studies and to shape a narrative that would deflect attention from Dupixent and redirect blame for the observed CTCL risk to an alternative culprit: atopic dermatitis itself. Defendants would also attempt to direct blame for CTCL to physicians despite prescribing Dupixent in accordance with the information and directions supplied by Defendants, and to further suggest that many CTCL diagnoses in Dupixent-treated patients were actually a benign, reversible reaction and not CTCL.

222. Defendants' employees and Defendants' retained KOLs submitted two letters to the editor of the Journal of the American Academy of Dermatology in response to the Hasan 2024 study, the first cohort study reporting a statistically significant increased risk of CTCL with Dupixent use.

223. The first letter was published on August 24, 2024 and co-authored by Yung-Tsu Cho, MD, PhD, who served as a sub-investigator for Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC and received speaker fees from Defendant, Sanofi-Aventis U.S. LLC, and Chia-Yu Chu, MD, PhD, who served as a sub-investigator for Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC, served as a consultant/advisory board member for Defendant, Sanofi-Aventis U.S. LLC, and received speaker fees from Defendant, Sanofi-Aventis U.S. LLC.¹⁸⁹ In this short letter, these authors argued that the atopic dermatitis population experiences an increased rate of CTCL and other lymphomas irrespective of Dupixent use and suggested it was possible that some proportion of reported CTCL cases in the Hasan study may actually have been lymphoid reactions rather than CTCL. Interestingly, the two publications

¹⁸⁹ Cho YT, Chu CY. Response to Hasan et al., "Dupilumab therapy for atopic dermatitis is associated with increased risk of cutaneous T-cell lymphoma: A retrospective cohort study". J Am Acad Dermatol. 2025;92(1):e9-e10. doi:10.1016/j.jaad.2024.06.105

cited by Cho and Chu in support of their claim that atopic dermatitis patients experience an elevated rate of CTCL were each co-authored by KOLs retained by Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC. They also made certain unfounded assumptions regarding the composition of the study groups without citing any data in support of their position, and ultimately advised fellow dermatologists to observe their patients for the development of certain adverse events rather than to exercise caution in prescribing Dupixent. Although no funding source was claimed, this letter was published Open Access at a price of \$4,200.

224. The second letter was funded by Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC and published on September 30, 2024.¹⁹⁰ This letter was co-authored by 10 employees of Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC, as follows: Tien V. Nguyen, MD, Andrea Fleisch Marcus, PhD, MPH, Stephane Levy, MD and Marius Ardeleanu, MD of Regeneron Pharmaceuticals Inc., Tarrytown, New York; Sarah-Jo Sinnott, MPharm, PhD, Debra Sierka, PharmD and Ana B. Rossi, MD, CMD of Sanofi, Cambridge, Massachusetts; Anna Coleman, MS of Regeneron Pharmaceuticals Inc., Dublin, Ireland; Carlos Peralta, MD, MBA of Sanofi, Amsterdam, Netherlands; and Prashant Kushwaha, MD of Sanofi, Mumbai, India. After performing what they referred to as a thoughtful evaluation of the study by Hasan and Colleagues, Defendants' employees swiftly proceeded to conclude the study was invalid and reached unsupported conclusions. In support of this position, Defendants' employees claimed that the atopic dermatitis population experiences an increased rate of CTCL, citing the same KOL-authored study relied upon by Cho and Chu and another industry-funded

¹⁹⁰ Nguyen TV, Marcus AF, Sinnott SJ, et al. Commentary: Response to "Dupilumab therapy for atopic dermatitis is associated with increased risk of cutaneous T cell lymphoma: A retrospective cohort study". J Am Acad Dermatol. 2025;92(2):e41-e42. doi:10.1016/j.jaad.2024.08.082

study also co-authored by one of Defendants' KOLs. They also claimed that the comparator group in the Hasan study was inappropriate, based upon their belief that those not treated with Dupixent were more likely to have less severe disease. Relatedly, they further advanced the unfounded assumption that "dupilumab is generally prescribed to patients with chronic, refractory AD", which runs contrary to the clinical treatment guidelines their employers funded that recommend Dupixent as a first-line treatment and advocate for its use in otherwise treatment-naïve patients. Finally, the authors stated "[a]s of March 28, 2024, ongoing safety surveillance activities for dupilumab have not identified an association with CTCL", and that "the available evidence does not support a causal association between dupilumab and CTCL".

225. Clearly, Defendants' employees took an untenable position as, upon information and belief, Defendants knew, or through the exercise of reasonable diligence should have known at the time this letter was first published that 4 retrospective cohort studies, 2 disproportionality analyses and no fewer than 20 case reports and case series encompassing dozens of patients had been published reporting an association between Dupixent and CTCL. This is, of course, in addition to the cases of CTCL observed during the Dupixent clinical development program, including the 3 cases reported in the long-term extension trial which published its results on ClinicalTrials.gov in October 2023 and in JAMA Dermatology in July 2024.

226. Defendants doubled down on their misrepresentations in a lengthy review of Dupixent safety data published in the American Journal of Clinical Dermatology in September 2025.¹⁹¹ This article was co-authored by 1 KOL (Richard G. Langley) and 10 employee-shareholders of Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi, as follows: Guy

¹⁹¹ Langley RG, Gherardi G, Coleman A, et al. The Safety Data of Dupilumab for the Treatment of Moderate-to-Severe Atopic Dermatitis in Infants, Children, Adolescents, and Adults. *Am J Clin Dermatol*. 2025;26(6):981-1002. doi:10.1007/s40257-025-00952-w

Gherardi of Sanofi, Reading, UK; Anna Coleman of Regeneron Pharmaceuticals Inc., Dublin, Ireland; Marius Ardeleanu, Stephane Levy, Ashish Bansal, Zhen Chen, Brad Shumel and Faisal A. Khokhar of Regeneron Pharmaceuticals Inc., Tarrytown, New York; Ainara Rodríguez-Marco of Sanofi, Madrid, Spain; and Ana B. Rossi, of Sanofi, Cambridge, Massachusetts. Although Defendants framed this review as being the “most comprehensive dupilumab safety assessment to date for the treatment of moderate-to-severe atopic dermatitis”, they instead used it as an opportunity to publicly misrepresent clinical trial data and the safety profile of Dupixent. As an initial matter, Defendants selectively omitted important safety data from the review by focusing only on a subset of the clinical trials that had been completed in the atopic dermatitis development program. Despite substantial evidence of an association between Dupixent and CTCL from 6 retrospective cohort studies, 5 disproportionality analyses and dozens of descriptive cohort studies, case reports, case series and systematic reviews – in addition to being notified by the FDA that the agency had identified a safety signal for CTCL with Dupixent – Defendants boldly, and falsely – and astonishingly, represented “ [t]here is currently no evidence associating CTCL and dupilumab treatment”. In alignment with their detraction playbook, Defendants then proceeded to blame atopic dermatitis for any reported CTCL risk, downplay the utility of spontaneous adverse event data analysis in postmarketing surveillance and discount any available research concerning Dupixent and CTCL, serially citing their own letter to the editor (Nguyen et al. 2025) and other pieces authored by company KOLs as support for their unsound position. Finally, Defendants utilized a full-page infographic to call attention to the findings of the review, highlighting, among other things, “No increased risk of...cutaneous T-cell lymphoma” with Dupixent.

227. Taking it a step further, two KOLs retained by Defendants, Emma Guttman-Yassky, MD, PhD and Patrick M. Brunner, MD, in conjunction with their colleagues, submitted

the Neubauer 2024 research letter discussed *supra* to the Journal of American Dermatology in response to the Hasan 2025 study which was published on September 7, 2024.¹⁹² Despite expressly claiming no funding sources in the publication, these authors included several statements indicative of strong bias and industry influence. Examples include “[w]e are concerned that this report might negatively influence physicians from prescribing dupilumab, causing more harm than good” and “[a]pproval of dupilumab has been life-changing for AD patients. Concerns over CTCL should not sway dermatologists from utilizing dupilumab for AD, with skin biopsy indicated in equivocal cases.” These statements reflect a conspicuous change in tone from the Owji 2023 study, also co-authored by Emma Guttman-Yassky, MD, PhD, which alternatively concluded that CTCL may “prove to be the exception” (to the lack of risk observed for other malignancies) and advised “Given the possible link between CTCL and dupilumab, additional studies evaluating this association are warranted”. Of note, during the 24-month period following publication of Owji 2023, Defendants compensated Emma Guttman-Yassky, MD, PhD \$130,715 in general payments and \$3,572 in research payments along with \$1,920,394 in associated research funding according to CMS OpenPayments data. Patrick M. Brunner, MD likewise received \$15,359 in general payments from Defendants during this period. Finally, despite declaring “no financial disclosures”, co-author Larisa J. Geskin, MD had also been paid \$29,610 in consulting fees by Regeneron Pharmaceuticals, Inc. in 2021 for work related to Dupixent.

228. Just 12 days prior to the publication of the Neubauer 2024 research letter, Emma Guttman-Yassky, MD, PhD and her colleagues submitted yet another study investigating malignancies among atopic dermatitis patients treated with Dupixent on August 25, 2024 to The

¹⁹² Neubauer ZJK, Brunner PM, Geskin LJ, Guttman E, Lipner SR. Decoupling the association of dupilumab with cutaneous T-cell lymphoma. J Am Acad Dermatol. 2024;91(6):1296-1298. doi:10.1016/j.jaad.2024.08.057

Journal of Allergy and Clinical Immunology: In Practice.¹⁹³ Using data from TriNetX, these authors ultimately reported that Dupixent-treated patients had a significantly lower risk of internal malignancy development compared with patients treated with other atopic dermatitis therapies. Oddly, however, these authors claimed there were not enough cases in TriNetX to examine CTCL as a specific endpoint, while nonetheless postulating that patients with severe atopic dermatitis may have a higher baseline lymphoma risk regardless of treatment exposure. Their claim of an inadequate number of CTCL cases is peculiar for several reasons, particularly given that somehow Emma Guttman-Yassky, MD, PhD and colleagues had an adequate number of cases to analyze CTCL as an endpoint when conducting almost the same exact study, at almost the same exact time, in the same database, yet followed patients over a period four times shorter (20 years vs. 5 years) in the Neubauer 2024 research letter. Also curious, senior author Nicholas Gulati, MD, PhD declared no conflicts of interest in the publication despite receiving 31 payments from Defendants, Regeneron Healthcare Solutions, Inc. and Genzyme Corporation in 2023 and 2024 according to CMS OpenPayments data.

229. One month after the publication of the Neubauer 2024 research letter, Patrick M. Brunner, MD, as senior author, submitted a review publication to The Journal of Allergy and Clinical Immunology self-citing his Neubauer research letter to bolster the perceived reliability and reach of the findings.¹⁹⁴ In this review, the authors cite the Hasan 2024 to simply note “[t]here have also been previous studies that associated an increased risk of Cutaneous T-cell lymphoma (CTCL) in patients treated with dupilumab for AD”, but without providing any of the results of

¹⁹³ Garate D, Thang CJ, Chang CT, et al. Risk of malignancy associated with use of dupilumab versus other treatments in atopic dermatitis patients: A national database analysis. *J Allergy Clin Immunol Pract.* 2025;13(3):698-701.e1. doi:10.1016/j.jaip.2024.11.015

¹⁹⁴ Meledathu S, Naidu MP, Brunner PM. Update on atopic dermatitis. *J Allergy Clin Immunol.* 2025;155(4):1124-1132. doi:10.1016/j.jaci.2025.01.013

the study. They also fail to cite the Owji 2023 study, Tsai 2024 study and Mandel 2024 study, all of which similarly identified an increased risk of CTCL in patients treated with Dupixent and had been published by this time. Brunner and his co-authors went on to describe in detail the findings of his research letter that was published the month prior, all while using the term “they” to refer to its authors in attempt to manufacture some perception of independence from the current work. Additionally, the authors made sure to mention that none of the cases of CTCL in the Lavin study reporting a safety signal for CTCL with Dupixent involved patients treated for non-cutaneous indications to suggest atopic dermatitis was driving the apparent CTCL risk. Notably, although Brunner and his co-authors discussed several classes of available therapies for the treatment of atopic dermatitis throughout their review, CTCL was only referenced in the section describing the only FDA-approved IL-4 receptor alpha inhibitor, Dupixent.

230. Initially submitted on December 31, 2024 and published February 13, 2025 in The Journal of Allergy and Clinical Immunology, an editorial authored by Defendants’ KOL Peck Y. Ong, MD sought downplay the safety signal for CTCL with Dupixent reported in the disproportionality analysis by Cabrera-Perez and colleagues.¹⁹⁵ In common refrain with other letters and editorials authored by Defendants’ KOLs, Ong carefully noted the Cabrera-Perez et al. “observation that CTCL was reported almost exclusively in patients with AD, but not in those with asthma or other atopic conditions, suggests that dupilumab by itself is not causative of CTCL”. Ong also advanced the theory that the CTCL cases in the Cabrera-Perez 2024 study may have actually been benign cutaneous lymphoid reactions that were misdiagnosed as CTCL, relying on another study authored by Defendants’ KOLs in support of this proposition. Despite acknowledging the findings of Cabrera-Perez et al. “support a worsening of CTCL with dupilumab

¹⁹⁵ Ong PY. Is dupilumab use in atopic dermatitis associated with cutaneous T-cell lymphoma?. J Allergy Clin Immunol. 2025;155(5):1481-1482. doi:10.1016/j.jaci.2025.02.002

use”, Ong boldly concluded “[t]here is no evidence that dupilumab causes or leads to the development of CTCL in AD”.

231. Another letter to the editor authored by multiple KOLs retained by Defendants aiming to allay concerns regarding the safety signal for CTCL with Dupixent reported by Lavin and colleagues was published in *The Journal of Investigative Dermatology* in June 2025.¹⁹⁶ These authors put forward an interesting theory that “[h]eightedened vigilance around CTCL shortly after the introduction of dupilumab for AD, may have led to over-reporting by dermatologists”, despite the absence of a warning regarding CTCL in the prescribing information for Dupixent or any communications from Defendants regarding a potential relationship between Dupixent and CTCL, in addition to the existence of the well-known phenomenon of significant underreporting of postmarketing adverse events. Although the disproportionality analysis by Lavin et al. covered a period during which Dupixent had been approved for 6 unique indications (5 indications other than atopic dermatitis) in the United States, the authors “suggest[ed] that comparison with drugs not commonly used for AD could be misleading”. Again, consistent with previous letters and editorials authored by Defendants’ KOLs, these authors commented “[n]one of the CTCL reports in this paper (or other literature) featured dupilumab for asthma”, further noting that an “[a]nalysis of associations in such people would be useful, as a lack of signal would further support the association being with AD over drug”. These authors additionally pronounced “reports of CTCL in patients with severe AD predate dupilumab and large-scale population-based evidence supports an increased risk of lymphoma in severe AD”. However, the only research they could find to support these claims included a 2020 cohort study which specifically excluded patients with CTCL

¹⁹⁶ James ML, Pink AE, Woolf RT, et al. Response to Lavin et al (2025)'s Paper: Cutaneous T-Cell Lymphoma after Dupilumab Use: A Real-World Pharmacovigilance Study of the FDA Adverse Event Reporting System. *J Invest Dermatol.* 2025;145(9):2325-2326. doi:10.1016/j.jid.2025.05.022

and a small 2004 case series of 4 patients with long-term atopic dermatitis who eventually developed lymphomatoid papulosis or cutaneous anaplastic large cell lymphoma. Ultimately, Defendants' KOLs concluded, "[w]e highlight that no clear causal association has been identified".

232. Next, craftily circumventing the hassles and uncertainties of the peer review process, Defendants leaned on their deep connections with members of scientific journal editorial and advisory boards to dispel CTCL concerns and champion the continued, unfettered use of Dupixent.

233. In September 2025, in response to the methodologically robust study by Sheng-Kai Ma and colleagues identifying a strong, significant association between Dupixent use and CTCL among asthma patients, Defendants' KOL Ian D. Pavord, DM wrote and published an editorial in the *European Respiratory Journal*.¹⁹⁷ Dr. Pavord, who is also a member of the International Advisory Board for the *European Respiratory Journal* and a contributor to Defendants' ADVENT platform, acknowledged "dupilumab may be associated with an increased risk of cutaneous lymphoma", while downplaying the magnitude of the observed risk as "very small", asserting that cutaneous lymphomas are "rarely life-threatening" and ultimately characterizing the findings of Sheng-Kai Ma et al (2025) as "much ado about nothing".

234. In October 2025, Tiago Torres, a KOL retained by Defendant, Genzyme Corporation ("Sanofi-Genzyme"), published an editorial in the *American Journal of Clinical Dermatology*, a journal for which he serves as an Editorial Board member, describing recent research concerning Dupixent and CTCL.¹⁹⁸ As with other pieces authored by Defendants and their KOLs, Dr. Torres casted Dupixent as having "revolutionized" atopic dermatitis treatment

¹⁹⁷ Davies TJ, Pavord ID. Lymphoma in patients with asthma treated with dupilumab: much ado about nothing?. *Eur Respir J*. 2025;66(3):2501321. Published 2025 Sep 25. doi:10.1183/13993003.01321-2025

¹⁹⁸ Torres T. Dupilumab and Cutaneous T-Cell Lymphoma: A Call for Vigilance, Not Alarm. *Am J Clin Dermatol*. Published online October 22, 2025. doi:10.1007/s40257-025-00991-3

while emphasizing that the risk of CTCL should not stop physicians from prescribing Dupixent, in triplicate, as follows: “overstating a malignancy risk could discourage clinicians from prescribing a highly effective and generally well-tolerated therapy, potentially depriving patients of substantial clinical benefit”; “concerns regarding CTCL in the context of dupilumab should not deter the use of a highly effective, safe, and well-tolerated therapy”; and “Does this mean dermatologists should be reluctant to prescribe dupilumab? Certainly not.” Nonetheless, Dr. Torres did concede in his editorial that the “CTCL signal” for Dupixent was “potentially real” while also describing various biological mechanisms providing “a plausible link between dupilumab and CTCL risk”.

235. Another Defendant-funded KOL and his colleagues argued against a causal link between Dupixent and CTCL in an editorial published in the *Journal of Drugs in Dermatology* in November 2025.¹⁹⁹ In their editorial, these authors repeatedly mischaracterized evidence concerning CTCL with Dupixent use, referred to findings from high quality epidemiologic studies and well-documented case series as “anecdotal”, and concluded that available research did not provide “convincing evidence of causality”. Despite acknowledging that causal mechanisms for Dupixent-induced CTCL that have been proposed in the literature “are plausible”, the authors suggested atopic dermatitis was more likely responsible for the observed CTCL risk, and ultimately advocated, “we do not believe dupilumab use should be limited due to concern about CTCL risk, particularly given its efficacy in treating chronic inflammatory diseases”. Curiously, while expressly declaring he “has no conflicts to disclose” in the disclosures section of this editorial, the senior author Adam Friedman, MD, FAAD has received over \$850,000 in general payments from Defendants, including \$519,143.97 from Defendant, Genzyme Corporation and

¹⁹⁹ Farah M, Zarabian N, Friedman A. Connecting the Plaques: Exploring the Link Between Dupilumab and Cutaneous T-cell Lymphoma. *J Drugs Dermatol*. 2025;24(11):1148-1149. doi:10.36849/JDD.1125

\$349,501.37 from Regeneron Healthcare Solutions, Inc., the subsidiary of Defendant, Regeneron Pharmaceuticals, Inc.

236. More recently, Defendants' efforts to manipulate the scientific record were exemplified in the Kridin 2026 study discussed *supra*. As noted, one of the co-authors of this study, Diamant Thaçi, is a consultant, speaker and/or advisory board member for Defendants. Additionally, the senior author of this study, Ralf J. Ludwig, is an editorial board member of the journal in which it was published. However, here the issue of conflict of interest extended beyond authorship, and even editorial board connections. Indeed, independent, unbiased peer review is a hallmark of academic publishing and is vital to ensure that published scholarly work reflects an honest and reliable application of the scientific process to advance knowledge on a given research question. Peer review is an essential gatekeeping function that maintains trust in the scientific research enterprise and safeguards against the publication of unwarranted claims, unacceptable interpretations or personal views without impartial appraisal and verification by experts in the field. Despite this imperative, interestingly, both of the peer reviewers charged with evaluating the rigor, integrity and veracity of this paper prior to its publication were also Defendants' KOLs. The first peer reviewer was Patrick Brunner, Defendants' KOL and co-author of the Neubauer 2024 research letter, and the second peer reviewer was Filip Rob, who also serves as a paid KOL for Defendants.²⁰⁰ Beyond design flaws and flouting scientific publishing conventions, other aspects of this publication are equally troubling. In their discussion section, the authors stated their findings were "in line with a recent report on no increased CTCL risk in AD patients", citing the Neubauer 2024 research letter co-authored by KOL – and now peer reviewer – Patrick Brunner.

²⁰⁰ See "Disclosure statement" in Rob F, Pinkova B, Sokolova K, Kopuleta J, Jiraskova Zakostelska Z, Cadova J. Real-world efficacy and safety of dupilumab in children with atopic dermatitis under 6 years of age: a retrospective multicentric study. *J Dermatolog Treat.* 2025;36(1):2460578. doi:10.1080/09546634.2025.2460578

They then went on to state that their findings “seemingly contrast[ed] with two previous reports on this topic”, citing only the Hasan 2024 study and the cross-sectional study by Hamp and colleagues. As a result, the authors framed the body of research concerning Dupixent and CTCL as consisting only of these four studies, with two finding an association and two finding no association. Such a discussion could hardly be considered balanced or accurately placing the present findings in appropriate context with extant research on the subject, and as such, this is something that unbiased peer reviewers would have demanded remedying prior to publication. Furthermore, in detailing their complaints with the Hasan 2024 study, these authors cited five different letters to the editor published in response to Hasan 2024, including the one by KOLs Cho and Chu, while omitting the letter from Nguyen and colleagues, employees of Defendants. Perhaps this omission was an attempt to distance the present work from Defendants’ influence. Finally, although the stated primary objective of the study was to clarify the association between atopic dermatitis and lymphoma, with evaluating the influence of Dupixent on lymphoma as a secondary aim, the authors chose to title their publication “Dupilumab treatment is not associated with changes in lymphoma risk in atopic dermatitis and other type 2 inflammatory diseases”.

237. After “observ[ing] increasing parental hesitancy about initiating dupilumab for their children” due to CTCL concerns, authors with no declared conflicts of interest published a review article in *Pediatric Dermatology* in May 2026 to characterize evidence and provide recommendations for counseling families.²⁰¹ As a threshold issue, many of the references in the paper do not support the proposition for which they were cited. It is unclear if these erroneous citations are simply a result of haphazard work or if they were intentional. Nonetheless, it is clear

²⁰¹ Bueth MG, Sy T, Eichenfield LF. Safety of Dupilumab and Risk of Cutaneous T-Cell Lymphoma in Pediatric Patients With Atopic Dermatitis: A Data-Driven Guide to Counseling Patients and Families. *Pediatr Dermatol*. 2026 (Ahead-of-Press). doi:10.1111/pde.70285

that supplying false and/or unsupported information to guide clinician conversations with families would be very concerning. These authors use their pages to push the narrative that the observed CTCL cases reflect “unmasking, not causation” as if it were a true, settled fact. They frame Dupixent as the “safest” option for children, with “alternatives carry[ing] more established risks” of lymphoma, also claiming that “dupilumab’s effects are reversible upon discontinuation”. Further, they suggest that a lack of reported CTCL cases in the exceptionally small (fewer than 200 patients) and non-diverse clinical trial population are reassuring, without acknowledging the lack of generalizability of these findings to real-world pediatric patients and omitting any reference to CTCL cases that have already been reported in real-world pediatric and adolescent patients treated with Dupixent. Despite declaring no conflicts of interest in their paper, the senior author, Lawrence F. Eichenfield received \$1,024,465.53 in general payments from Defendants and their subsidiaries between 2019 and 2025 according to CMS OpenPayments data.

238. Defendants provided direct funding to four KOLs to write another letter to the editor in response to the Sheng-Kai Ma 2025 study in an effort to protect the Dupixent brand and ensure that the study findings did not “cause detrimental impacts on prescriber practices”.²⁰² This letter was published in European Respiratory Journal in July 2026. While prominently declaring “crucial methodological flaws, biological implausibility, and incident risk of disease focus” served to invalidate the study, the authors’ farrago of critiques instead suggests a lack of familiarity with the TriNetX platform and epidemiologic research generally, with certain comments bringing to question whether they actually read the study. For example, because analytical cohort studies are designed to measure and compare risk (also referred to as incidence) of disease among exposed

²⁰² Bacharier LB, Busse WW, Jackson DJ, Katial RK. Critical flaws in: "Dupilumab and lymphoma risk among patients with asthma: a population-based cohort study". Eur Respir J. 2026;68(1):2501565. doi:10.1183/13993003.01565-2025

and unexposed populations, it is unclear how an “incident risk of disease focus” could possibly be considered a “flaw” or “area[] of concern” of the Sheng-Kai Ma 2025 study. Oddly, despite concluding that the study had “considerable limitations and flaws”, these authors somehow reasoned that the finding of lower all-cause mortality was not subject to these apparent shortcomings. Of note, these four KOLs collectively received \$411,937.20 in general payments from Defendants and their subsidiaries in 2025 alone according to CMS OpenPayments data.

239. Beyond publication, Defendants also leaned on their KOLs and relationships with patient advocacy organizations to relay misleading messaging regarding Dupixent and CTCL risk, assuage patient concerns and ensure continued utilization of Dupixent through podcasts and webpages.^{203,204}

240. As reflected above, Defendants dedicated considerable time, effort and financial resources to shaping an elaborate narrative that would downplay the risk of CTCL with Dupixent through publications and other mediums. At no point did Defendants acknowledge the merits of the findings produced through independent research or take initiative to commit to conducting necessary studies to better understand the pathological mechanisms of Dupixent-induced CTCL or discover why Dupixent triggers rapid and severe progression of subclinical CTCL.

H. Defendants’ Labeling and Instructions for Dupixent

241. From its original approval in 2017 to the present, the United States prescribing information for Dupixent has been deficient in that it has failed to provide adequate instructions

²⁰³ Sanofi Campus. DermTalks Podcast: Atopic Dermatitis in Special Populations – CTCL February 19, 2025. Last accessed July 27, 2026. <https://www.campus.sanofi/se/hidden-pages/ar/atopic-dermatitis-cutaneous-t-cell-lymphoma> (“Based on prior evaluations and ongoing safety surveillance, the available evidence does not support a causal association between dupilumab and CTCL”...“There is no strong scientific evidence regarding the relationship of CTCL occurrence and dupilumab treated patients.”)

²⁰⁴ Eczema Support Australia. Dupilumab (Dupixent®) and Cutaneous T-Cell Lymphoma (CTCL). April 17, 2026. Last accessed July 27, 2026. <https://www.eczemasupport.org.au/dupilumab-dupixent-and-cutaneous-t-cell-lymphoma-ctcl/> (“No causal link has been found”...“The bottom line: No research has proven that dupilumab causes skin lymphoma.”)

for its safe prescription and use. At all times relevant hereto, the United States prescribing information for Dupixent has contained **1)** no BOXED WARNING advising patients and their healthcare providers of the risk of CTCL or any other form of cancer with Dupixent treatment; **2)** no warnings in section 5, WARNINGS AND PRECAUTIONS, advising patients and their healthcare providers of the risk of CTCL or any other form of cancer with Dupixent treatment; **3)** no reference to CTCL in section 6, ADVERSE REACTIONS, advising patients and their healthcare providers that clinically significant CTCL adverse reactions have been reported with the use of Dupixent; **4)** no contraindication for use in patients with a history of CTCL or active CTCL; **5)** no instructions directing healthcare providers to perform adequate testing of patients (through biopsy or other means) to confirm a clinical diagnosis of atopic dermatitis, particularly in equivocal cases, before commencing treatment with Dupixent; **6)** no instructions directing healthcare providers to perform adequate testing of patients (through biopsy, cytometry or other means) to rule out the presence of subclinical or smoldering CTCL before commencing treatment with Dupixent; **7)** no instructions directing healthcare providers to perform regular and continuous monitoring during treatment with Dupixent for the development of signs and symptoms of CTCL and repeat biopsies to detect changes in pathologies during treatment with Dupixent consistent with CTCL development; and **8)** no instructions directing healthcare providers to discontinue use of Dupixent when a patient exhibits signs of non-responsiveness to treatment or worsening of disease, or otherwise develops signs and symptoms consistent with CTCL.

242. Defendants' failure to add the above-referenced information to the United States prescribing information for Dupixent is despite sufficient notice of reasonable evidence of a causal association between Dupixent use and incident CTCL that would have supported inclusion of this information well before Plaintiff's use of Dupixent.

243. Defendants' failure to appropriately revise the United States prescribing information for Dupixent to reflect the clinically important risk of CTCL with Dupixent treatment continued in the face of mounting evidence from dozens and dozens of postmarketing adverse events, published case reports, published case series, descriptive observational studies, analytic cohort studies and disproportionality analyses.

244. The vast body of scientific evidence discussed *supra* establishes that there exists a strong and consistent causal relationship between the use of Dupixent and incident CTCL, to include both new cases of CTCL and rapid exacerbation of subclinical CTCL, which was known or knowable to Defendants at all times relevant hereto. Further, at all times relevant hereto, numerous practical measures to prevent, mitigate and/or reduce the risk of Dupixent-induced CTCL were known or knowable to Defendants.

245. At all times relevant hereto, Defendants maintained complete responsibility for the content of the United States Prescribing Information for Dupixent.

246. At all times relevant hereto, Defendants were under a duty to advise healthcare providers within Section 5, WARNINGS AND PRECAUTIONS of the United States Prescribing Information for Dupixent, of clinically significant adverse reactions, including any that are potentially fatal, that are serious even if infrequent or that can be prevented or mitigated through appropriate use of the drug, along with limitations in use imposed by them, associated with the use of Dupixent. (21 CFR 201.57(c)(6)(i)).

247. At all times relevant hereto, Defendants were under a duty to advise healthcare providers within Section 5, WARNINGS AND PRECAUTIONS of the United States Prescribing Information for Dupixent, of laboratory tests that should be performed before, during and after therapy to follow patient responses or identify possible adverse reactions associated with the use

of Dupixent. (21 CFR 201.57(c)(6)(iii)).

248. At all times relevant hereto, Defendants were under a duty to advise healthcare providers within a BOXED WARNING in the United States Prescribing Information for Dupixent, of clinically significant adverse reactions, particularly those that may lead to death or serious injury, associated with the use of Dupixent. (21 CFR 201.57(c)(1)).

249. At all times relevant hereto, Defendants were under a duty to promptly revise the United States Prescribing Information for Dupixent to add warnings regarding clinically significant hazards as soon as there is reasonable evidence of a causal association with the drug. (21 CFR 201.57(c)(6)(i)). Importantly, however, a causal relationship need not have been definitely established in order to add such a warning to the United States Prescribing Information.

250. At all times relevant hereto, through submitting to the FDA a Changes Being Effected (CBE) Supplement, Defendants had available to them the option to unilaterally add new warnings regarding the risk of CTCL to the United States Prescribing Information for Dupixent reflecting newly-acquired information during the postmarketing period evidencing this clinically significant risk. (21 CFR 314.70(c)).

251. Similarly, through submitting a CBE Supplement to the FDA, Defendants had available to them the option to add new contraindications for use of Dupixent in patients without histopathologically confirmed atopic dermatitis or at risk of developing or experiencing exacerbation of subclinical CTCL and/or to add or strengthen instructions concerning dosage and administration of Dupixent that would serve to increase the safety of its use. (21 CFR 314.70(c))

252. Notwithstanding, Defendants chose not to undertake any such actions.

253. Further, at all times relevant hereto, Defendants failed to take other reasonable and prudent steps within their capacity, including, but not limited to sending “Dear Healthcare

Provider” (DHCP) letters or issuing and disseminating safety alerts, to advise patients and healthcare providers that Dupixent treatment carries a significant risk of CTCL and to convey appropriate mitigation strategies to reduce or eliminate the risk of Dupixent-induced CTCL. Similarly, Defendants failed to advise patients and healthcare providers of new research evidencing a significant risk of CTCL with Dupixent use through prudent and reasonable means within their capacity.

I. FDA Identifies Safety Signal for CTCL with Dupixent

254. On January 15, 2025, the FDA notified Defendant, Regeneron Pharmaceuticals, Inc. via email that it identified and began evaluating a newly identified safety signal (NISS) regarding “dupilumab therapy for atopic dermatitis being associated with increased risk of cutaneous T cell lymphoma” on December 26, 2024. The FDA further noted that the agency had classified this NISS as a potential risk and that it was informing Regeneron Pharmaceuticals, Inc. in accordance with the Food and Drug Administration Amendments Act of 2007, the Food Drug & Cosmetic Act and their implementing regulations.^{205,206}

255. On March 31, 2025, the FDA published its quarterly report “Potential Signals of Serious Risks/New Safety Information Identified by the FDA Adverse Event Reporting System (FAERS)” for the 4th quarter of 2024.²⁰⁷ In this report, FDA identified CTCL as a potential signal of a serious risk associated with Dupixent, and noted it was “evaluating the need for regulatory

²⁰⁵ Center for Drug Evaluation and Research. Manual of Policies and Procedures: MAPP 6700.9: FDA Posting of Potential Signals of Serious Risks Identified by the FDA Adverse Event Reporting System. U.S. Food and Drug Administration. September 9, 2019.

²⁰⁶ Center for Biologics Evaluation and Research. Standard Operating Procedures and Policies: SOPP 8420: FDAAA Section 921: Posting of Potential Signals of Serious Risk. U.S. Food and Drug Administration. February 27, 2022.

²⁰⁷ October - December 2024 | Potential Signals of Serious Risks/New Safety Information Identified by the FDA Adverse Event Reporting System (FAERS). U.S. Food and Drug Administration. March 31, 2025. <https://www.fda.gov/drugs/fdas-adverse-event-reporting-system-faers/october-december-2024-potential-signals-serious-risksnew-safety-information-identified-fda-adverse>

action”.

256. As of the filing of this complaint, Defendants have still failed to implement appropriate labeling revisions to add necessary information regarding the risk of CTCL with Dupixent use or appropriate mitigation strategies to reduce or eliminate the risk of Dupixent-induced CTCL.

FACTS SPECIFIC TO PLAINTIFF

257. In February 2021, Plaintiff, Anne Gibson was prescribed Dupixent to treat her atopic dermatitis.

258. Plaintiff, Anne Gibson used Dupixent in a manner consistent with and as intended by Defendants.

259. In total, Plaintiff, Anne Gibson used Dupixent from March 2021 to February 2023.

260. In or about February 2023, Plaintiff, Anne Gibson was diagnosed with mycosis fungoides.

261. As a result of Plaintiff's CTCL diagnosis, she has required extensive treatments and the use of medications including phototherapy, chemotherapy, and radiation.

262. Plaintiff, Anne Gibson has been advised by her healthcare providers that she will likely require ongoing treatments and healthcare services for CTCL indefinitely.

263. Before prescribing and using Dupixent, Plaintiff, Anne Gibson and Plaintiff's healthcare providers were exposed to Defendants' branded marketing for Dupixent and unbranded disease state awareness TV campaigns.

264. But for Defendants' failure to provide a clear warning regarding the increased risk of CTCL with the use of Dupixent, Plaintiff's physician would not have prescribed her Dupixent and Plaintiff, Anne Gibson would not have taken Dupixent to treat her atopic dermatitis.

265. At the time Plaintiff, Anne Gibson was first prescribed Dupixent, the available scientific data were adequate to put Defendants on notice that there existed reasonable evidence of causal association between Dupixent and development and progression of CTCL.

266. Throughout the relevant period, Plaintiff exercised reasonable diligence by seeking medical care, undergoing diagnostic testing, and following the advice of her treating physicians.

267. Plaintiff reasonably relied on Defendants' representations and omissions, the absence of warnings in Dupixent's prescribing information or any other publicly disseminated information in not connecting her CTCL diagnosis with Dupixent until after the public dissemination of information related to the FDA's posting of a newly identified potential safety signal (NISS) on March 31, 2025.

268. As a direct and proximate result of the dangerous and defective nature of Dupixent as described herein, Plaintiff suffered severe bodily injury, and resulting pain and suffering, disability, disfigurement, mental anguish, loss of capacity for the enjoyment of life, and expense of hospitalization and treatment. The losses are permanent and Plaintiff will continue to suffer the losses in the future.

FIRST CAUSE OF ACTION
STRICT LIABILITY – FAILURE TO WARN

269. Plaintiff incorporates by reference herein each of the allegations heretofore set forth in this Complaint as though fully set forth herein.

270. At all times relevant hereto, Dupixent was defective and unreasonably dangerous when it left the possession of Defendants in that it failed to contain warnings of an adequate or sufficient nature as to alert consumers and physicians, including Plaintiff and Plaintiff's healthcare providers, to the dangerous risks associated therewith, including, but not limited, to its propensity to cause serious and permanent injuries including those which Plaintiff sustained. These risks and

dangers were known and/or reasonably knowable by Defendants prior to and during the time which Plaintiff was prescribed and ingested Dupixent.

271. At all times relevant hereto, Defendants failed to provide sufficient warnings and instructions that would have put the general public, including Plaintiff and Plaintiff's physicians, on notice of the dangers and adverse effects caused by administration of Dupixent, including, but not limited to CTCL.

272. Defendants failed to provide warnings of such risks and dangers to Plaintiff and Plaintiff's healthcare providers as described herein, and further, concealed the known risks and dangers and failed to warn of known or scientifically knowable risks and dangers associated with Dupixent from patients, the medical community, and consumers, including Plaintiff, and Plaintiff's healthcare providers.

273. Plaintiff was prescribed and did administer Defendants' Dupixent in a manner consistent with and as intended by Defendants.

274. Ordinary patients and consumers, such as Plaintiff, could not have discovered or recognized any relevant potential risks and dangerous defects in Defendants' Dupixent through the exercise of reasonable care within their capacity.

275. Defendants, as entities materially involved in the development, testing, manufacture, sale and/or distribution of Dupixent, are held to the level of knowledge of an expert in the field.

276. Plaintiff individually, and through her prescribing physician, reasonably relied upon the skill, superior knowledge, and judgment of Defendants.

277. Despite their possession of knowledge regarding this risk and their duty to adequately warn of severe and dangerous adverse events associated with use of Dupixent,

Defendants failed to properly warn the medical community and consumers, including Plaintiff and Plaintiff's healthcare providers, that use of Dupixent was associated with an increased risk of CTCL.

278. Dupixent was designed, manufactured, distributed, sold and/or supplied by Defendants, and was marketed while defective due to inadequate warnings, instructions, labeling and/or inadequate testing in light of Defendants' knowledge of Dupixent's innate risks and dangers and attributable CTCL adverse events.

279. As a direct and proximate result of the dangerous and defective nature of Dupixent, Plaintiff suffered mycosis fungoides, and resulting pain and suffering, disability, disfigurement, mental anguish, loss of capacity for the enjoyment of life, expense of hospitalization, and medical treatment. These losses are permanent and the Plaintiff will continue to suffer the losses in the future.

SECOND CAUSE OF ACTION **NEGLIGENCE**

280. Plaintiff incorporates by reference herein each of the allegations heretofore set forth in this Complaint as though fully set forth herein.

281. At all times relevant hereto, Defendants were under a duty to exercise reasonable care in advertising, analyzing, assembling, compounding, designing, developing, distributing, formulating, inspecting, labeling, manufacturing, marketing, packing, producing, promoting, processing, researching, selling, and testing Dupixent to ensure that use of Dupixent did not result in avoidable injuries.

282. At all times relevant hereto, Defendants owed a duty to consumers, physicians, and the general public to assess, manage, and communicate the risks, dangers, and adverse effects of Dupixent, and to warn consumers and the medical community, including Plaintiff and Plaintiff's

healthcare providers, of those risks, dangers, and adverse effects.

283. Defendants' duties include, but are not limited to, carefully and properly advertising, analyzing, assembling, compounding, designing, developing, distributing, formulating, inspecting, labeling, manufacturing, marketing, packing, producing, promoting, processing, researching, selling, and testing Dupixent, which was placed in the stream of commerce, and providing adequate information regarding the appropriate use of Dupixent.

284. Defendants negligently breached the above-described duties to Plaintiff by committing negligent acts and/or omissions, including, but not limited to the following:

- a. failing in their obligation to provide consumers and the medical community, including Plaintiff and Plaintiff's healthcare providers, with accurate, adequate and clinically relevant information, data and warnings regarding the risk of CTCL associated with use of Dupixent, and/or that there existed safer alternative pharmaceutical drugs to treat atopic dermatitis;
- b. failing to continually monitor, test, and analyze data regarding safety, efficacy, and the prescribing practices for Dupixent;
- c. failing to review all adverse drug event information and to report any information bearing upon the adequacy and/or accuracy of its warnings, efficacy, or safety, including the risks and/or prevalence of side effects, including CTCL, caused by Dupixent to consumers and the medical community, including Plaintiff and Plaintiff's healthcare providers;
- d. failing to provide adequate post-marketing warnings and instructions after Defendants knew or should have known of the significant risks of, among other things, serious adverse events and/or reactions, including, but not limited to an increased risk of CTCL associated with use of Dupixent;
- e. failing to review all medical literature regarding Dupixent and failing to report data regarding the adequacy and/or accuracy of its warnings, efficacy, or safety of Dupixent specifically as it relates to CTCL;
- f. failing to disclose the results of the testing and other information in their possession regarding the potential for Dupixent to cause serious adverse events including, but not limited to, an increased risk of CTCL;
- g. representing that Dupixent was safe for use when, in fact, Defendants knew or should have known that it was unsafe for use and that Dupixent use was

associated with serious adverse events and/or reactions, including, but not limited to an increased risk of CTCL;

- h. promoting and marketing Dupixent for use despite the fact that Defendants knew or should have known that Dupixent use was associated with serious adverse events and/or reactions, including, but not limited to an increased risk of CTCL;
- i. promoting and marketing Dupixent as safe and effective for use when, in fact, it was unsafe, especially as compared to other available therapies to treat atopic dermatitis;
- j. failing to act as reasonably prudent drug manufacturers in advertising, analyzing, assembling, compounding, designing, developing, distributing, formulating, inspecting, labeling, manufacturing, marketing, packaging, producing, promoting, processing, researching, selling, and testing Dupixent;
- k. failing to exercise ordinary care in advertising, analyzing, assembling, compounding, designing, developing, distributing, formulating, inspecting, labeling, manufacturing, marketing, packing, producing, promoting, processing, researching, selling, and testing Dupixent so as to reveal and communicate the risk of serious adverse events and/or reactions, including, but not limited to, an increased risk of CTCL to consumers and the medical community, including Plaintiff and Plaintiff's healthcare providers;
- l. failing to conduct adequate post-marketing studies, non-clinical and clinical testing, and post-marketing surveillance and analyses to determine and subsequently communicate the safety profile and side effects associated with the use of Dupixent;
- m. continuing to promote the safety and effectiveness of Dupixent while downplaying its risks, even after Defendants knew or should have known of the significant risks of Dupixent use;
- n. failing to provide consumers and the medical community, including Plaintiff and Plaintiff's healthcare providers, with scientific data which indicated that Dupixent was unreasonably dangerous due to its propensity to cause serious adverse events including, but not limited to, an increased risk of CTCL;
- o. negligently and carelessly over-promoting Dupixent in a zealous and unreasonable manner, without regard for the potential dangers which it posed to users; and/or
- p. failing to adequately test Dupixent on patients typical of the atopic dermatitis population especially in light of the plan to market precisely

to that population of patients.

285. Although Defendants knew or should have known that Dupixent causes unreasonably dangerous side effects, including an increased risk of CTCL, they continue to market Dupixent, despite the fact there are safer and more or equally effective alternative therapies to treat atopic dermatitis.

286. Defendants knew or should have known that failure to exercise ordinary care, as described herein, would result in serious injury to patients, such as Plaintiff.

287. As a direct and proximate result of the dangerous and defective nature of Dupixent Plaintiff suffered mycosis fungoides, and resulting pain and suffering, disability, disfigurement, mental anguish, loss of capacity for the enjoyment of life, expense of hospitalization, and medical care and treatment. These losses are permanent and the Plaintiff will continue to suffer the losses in the future.

THIRD CAUSE OF ACTION **FRAUDULENT MISREPRESENTATION**

288. Plaintiff incorporates by reference herein each of the allegations heretofore set forth in this Complaint as though fully set forth herein.

289. Defendants' fraudulent, intentional and material misrepresentations and omissions regarding the safety and efficacy of Dupixent and of Dupixent's side effects, including that concerning an increased risk of CTCL, were communicated to Plaintiff directly through promotional materials, advertising, package inserts, journal publications, and the product monograph with the intent that Plaintiff use Dupixent. The safety and efficacy of Dupixent was also fraudulently and intentionally misrepresented to Plaintiff's healthcare providers with the intent that such misrepresentations would result in Dupixent being prescribed and administered to Plaintiff.

290. Defendants knew that the material representations they were making regarding the safety, efficacy and side effects of Dupixent were false.

291. Defendants fraudulently and intentionally made misrepresentations and/or actively concealed, suppressed, or omitted this material information with the intention and specific desire to induce consumers and the medical community, including Plaintiff and Plaintiff's healthcare providers, to use, prescribe and purchase Dupixent.

292. Defendants fraudulently and intentionally knew that Plaintiff and/or Plaintiff's healthcare providers would rely upon such material misrepresentations and/or omissions in selecting Dupixent for the treatment of Plaintiff.

293. Plaintiff and Plaintiff's healthcare providers detrimentally relied on the fraudulent misrepresentations outlined below.

294. Defendants made these material misrepresentations and/or omissions and actively concealed adverse information at a time when they, their agents and/or their employees knew that Dupixent had certain defects, dangers and characteristics that differed from what had been represented to the medical community and the consuming public, including Plaintiff's healthcare providers and Plaintiff.

295. Defendants misrepresented critical aspects of Dupixent's development and safety evaluation, including its carcinogenic risks and the scope of its clinical testing. As intended, these misrepresentations induced reliance of health care providers and drove prescriptions of Dupixent including Plaintiff's Dupixent prescription.

- a. Defendants publicly portray Dupixent as safe, thereby implying that they have adequately evaluated its carcinogenic potential, despite never conducted formal studies to assess mechanisms by which Dupixent may cause CTCL or accelerate its progression;
- b. Defendants misrepresented the results of premarket clinical trials and postmarketing observational studies evidencing a serious risk of CTCL with

the use of Dupixent; and

- c. Defendants promoted Dupixent as a safe and effective treatment for patients lacking diagnostic confirmation of atopic dermatitis and other inflammatory conditions despite the fact that the Defendants had not properly studied Dupixent in that patient population.

296. Defendants partnered, funded, and leveraged influential “third parties” to misrepresent the safety and effectiveness of Dupixent and promote its use in an unrestricted manner inconsistent with its FDA-approved indications. Defendants’ fraudulent conduct, as outlined below, led to the treatment guidelines health care providers rely upon to prescribe Dupixent to patients, including Plaintiff.

- a. Through a 2017 “steering committee” funded by Defendants, which later published in The Journal of Allergy and Clinical Immunology: In Practice, Defendants falsely characterized Dupixent’s safety profile as being superior to existing therapies, including in terms of lymphoma risk, notwithstanding that it had not yet been used by any patients under real-world conditions;
- b. Defendants used this self-funded publication within their treatment guidelines, falsely recommending Dupixent as a first-line systemic treatment option in adults with moderate-to-severe AD despite the fact that: (1) Dupixent was not FDA-approved for any indication at the time the steering committee met or formulated its recommendations, and (2) Dupixent has never been approved by the FDA as a first-line systemic therapy for atopic dermatitis; and
- c. Defendants fraudulently represented this “steering committee” recommendations and manuscript were free from bias, despite the fact that 10 of the 11 authors of the guidelines article being retained by Defendants as physician KOLs who had been paid a total of \$919,595 for consulting fees, food and beverage costs, education costs and travel and lodging expenses related to Dupixent.

297. Defendants, their agents and/or their employees, authored and funded misleading publications in the medical literature that contained false material facts and suppressed the truth regarding Dupixent’s safety, efficiency, and association with CTCL, to which Defendants knew or should have known.

- a. Upon information and belief, and as discovery is expected to reveal, authors

Yung-Tsu Cho and Chia-Yu Chu, acting at the direction of and in coordination with Defendants, published an August 2024 letter that sought to minimize the growing evidence linking Dupixent to CTCL, the precise injury suffered by Plaintiff, while failing to disclose Defendants' funding, involvement, and/or influence over the publication. This misleading communication was disseminated despite Defendants' knowledge of CTCL cases observed during the Dupixent clinical development program including at least three CTCL cases reported in the long-term extension trial.

- b. Despite there being 4 retrospective cohort studies, 2 disproportionality analyses and no fewer than 20 case reports and case series encompassing dozens of patients that had been published reporting an association between Dupixent and CTCL, on September 30, 2024, 10 employees of Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC published a letter misrepresenting material scientific evidence concerning the rate of CTCL among the atopic dermatitis population and their knowledge of the existence of a safety signal for CTCL with Dupixent, omitting that their marketing targeted patients with less severe forms of atopic dermatitis – the opposite of the population Defendants asserted was likely to comprise the Dupixent treated group in the Hasan 2024 study;
- c. Defendants further perpetrated fraudulent misrepresentations and omissions by coordinating a series of publications authored by their retained and compensated key opinion leaders, including the Neubauer 2024 research letter, a contemporaneous TriNetX analysis, and a subsequent review article, that falsely minimized the Dupixent–CTCL safety signal, selectively omitted contrary data, misrepresented study findings and comparator groups, and concealed significant financial conflicts while urging clinicians not to allow CTCL evidence to influence their prescribing decisions;
- d. In September 2025, Defendants falsely represented that there is no evidence supporting an association between CTCL and Dupixent, while being aware of at least 6 retrospective cohort studies, 5 disproportionality analyses and dozens of descriptive cohort studies, case reports, case series and systematic reviews. In addition, at that time, Defendants had already been notified by the FDA that the agency had identified a safety signal for CTCL with Dupixent;
- e. A published letter in June 2025 by Defendants' retained KOLs falsely suggested that heightened clinician “vigilance” led to over-reporting of CTCL with Dupixent, despite the well-known phenomenon of significant underreporting of postmarketing adverse events, and falsely asserting, without support, that no clear causal association had been identified;
- f. Defendants further advanced their fraudulent misrepresentations and omissions through a coordinated series of 2025 publications authored by

their retained and heavily compensated KOLs, including Ian D. Pavord, Tiago Torres, and Adam Friedman, MD, which collectively downplayed the strength of emerging epidemiologic evidence linking Dupixent to CTCL, mischaracterized high-quality studies as “anecdotal,” minimized the magnitude and seriousness of observed risks, urged clinicians not to let CTCL concerns deter prescribing, selectively emphasized Dupixent’s purported benefits while concealing and misrepresenting causal mechanisms, and failed to disclose substantial financial conflicts despite publicly claiming none existed;

- g. In January 2026, through their retained KOLs, Defendants mischaracterized results of the Kridin 2026 study and Talasila 2026 research letter as reassuring evidence of Dupixent’s safety by portraying the findings as reflecting a “decreased risk association,” “partial amelioration” of lymphoma risk, or mere “diagnostic confounding and delayed recognition”, despite conversely identifying an increased risk of CTCL with Dupixent in multiple populations, consistent with other studies reporting statistically significant increased risks of lymphoma among Dupixent-treated patients, while further providing no reliable basis to exclude a causal association with CTCL and mischaracterizing and/or omitting other available evidence relevant to the Dupixent-CTCL relationship.

298. Despite having a wealth of knowledge, since 2017, Defendants have at all times failed to disclose that the prescribing information for Dupixent is materially deficient. Each of the following failures constitute a fraudulent omission of safety information necessary for patients and prescribers, including Plaintiff, to be informed on the prescription and use of Dupixent:

- a. failure to include a BOXED WARNING advising patients and their healthcare providers of the risk of CTCL or any other form of cancer with Dupixent treatment;
- b. failure to include warnings in section 5, WARNINGS AND PRECAUTIONS, advising patients and their healthcare providers of the risk of CTCL or any other form of cancer with Dupixent treatment;
- c. failure to reference to CTCL in section 6, ADVERSE REACTIONS, advising patients and their healthcare providers that clinically significant CTCL adverse reactions have been reported with the use of Dupixent;
- d. failure to include a contraindication for use in patients with a history of CTCL or active CTCL;
- e. failure to include instructions directing healthcare providers to perform adequate testing of patients (through biopsy or other means) to confirm a

clinical diagnosis of atopic dermatitis, particularly in equivocal cases, before commencing treatment with Dupixent;

- f. failure to include instructions directing healthcare providers to perform adequate testing of patients (through biopsy, cytometry or other means) to rule out the presence of subclinical or smoldering CTCL before commencing treatment with Dupixent;
- g. failure to include instructions directing healthcare providers to perform regular and continuous monitoring during treatment with Dupixent for the development of signs and symptoms of CTCL and repeat biopsies to detect changes in pathologies during treatment with Dupixent consistent with CTCL development; and
- h. failure to include instructions directing healthcare providers to discontinue use of Dupixent when a patient exhibits signs of non-responsiveness to treatment or worsening of disease, or otherwise develops signs and symptoms consistent with CTCL.

299. Additionally, Defendants affirmatively highlighted the absence of a boxed warning and the lack of any requirement for initial laboratory testing or ongoing laboratory monitoring as favorable safety attributes of Dupixent in their promotional and patient-facing materials. In doing so, Defendants mislead patients and prescribing physicians that Dupixent posed no serious or latent safety risks and had been adequately evaluated for long-term adverse outcomes.

300. To drive profits, Defendants intentionally misrepresented and omitted material information through a multifaceted marketing strategy, spanning direct-to-consumer advertising, disease-awareness initiatives, and other promotional mechanisms. The following marketing strategies, executed by Defendants, their agents and/or their employees, were relied on by Plaintiff and her prescribing physician:

- a. Defendants misrepresented Dupixent's safety and effectiveness through provocative direct-to-consumer advertising that used glossy imagery and slogans such as "Du-More," "Better Days," and "This is Better" to falsely imply that patients would achieve materially happier, healthier, and more fulfilling lives by using Dupixent;
- b. Defendants further executed fraudulent misrepresentations and omissions by deploying the Dupixent MyWay and MyWay Ambassador programs,

related smartphone applications, and a network of unbranded disease-state awareness campaigns and websites, including UnderstandAD.com, EczemaExposed.com, ThinkCOPDInflammation.com, and LiveWithCOPD.com, to create the false impression of neutral educational resources while in fact steering patients to view their symptoms as serious and uncontrolled, encouraging “productive” doctor conversations designed to yield Dupixent prescriptions, and seeding demand for Dupixent even before FDA approval, all while concealing that these materials were crafted and controlled by Defendants to promote Dupixent’s use;

- c. Defendants fraudulently misrepresented diagnostic tools, including the Atopic Dermatitis Control Tool (ADCT) and related pediatric itch scales neutral clinical instruments, when in fact their high sensitivity artificially inflated atopic dermatitis diagnoses and artificially steered patients and physicians toward Dupixent treatment; and
- d. Through continuing medical education programs and their ADVENT global medical education platform, Defendants misrepresented promotional materials as unbiased scientific education, concealing their true marketing intent to increase Dupixent prescribing.

301. At no point did Defendants acknowledge the merits of the findings produced through independent research or take initiative to commit to conducting necessary studies to better understand the pathological mechanisms of Dupixent-induced CTCL or discover why Dupixent triggers rapid and severe progression of subclinical CTCL, instead Defendants misrepresented material facts to the public by funding and influence.

302. As further detailed above, Defendants engaged in the following fraudulent practices designed to induce healthcare providers and consumers to prescribe and use Dupixent, which were relied on by Plaintiff’s prescribing physician:

- a. inducing healthcare providers to switch their patients from other medications to Dupixent despite lacking the necessary evidence to demonstrate that Dupixent is safe and effective in this population and despite affirmative evidence that the drug is not safe in this patient population;
- b. inducing healthcare providers, through marketing efforts, to prescribe Dupixent to their patients without first obtaining diagnostic confirmation of an underlying indicated health condition;
- c. engaging in marketing efforts, such as disease state awareness campaigns

and branded advertising campaigns, seeking to induce patients to obtain a diagnosis for an indicated health condition and then specifically request a prescription of Dupixent from their healthcare provider when its use is not necessary; and

- d. intentionally misleading healthcare providers and the general public by making superiority claims for Dupixent as compared to other medications despite possessing the knowledge that these claims are false.

303. Defendants' misrepresentations were widely disseminated to healthcare providers through CME programs, clinical guidelines, KOL publications, and disease awareness campaigns, and to consumers via direct-to-consumer advertising, MyWay materials, and unbranded disease-state websites. These misrepresentations reached Plaintiff and Plaintiff's treating providers and directly influenced the decision to prescribe and use Dupixent.

304. Defendants' material misrepresentations and/or active concealment, suppression, and omissions were perpetuated directly and/or indirectly by Defendants, their sales representatives, employees, distributors, agents and/or detail persons, through databases, printouts, monographs, product labeling and other information drafted, prepared, marketed, sold, and supplied by Defendants, their sales representatives, employees, distributors, agents and/or detail persons.

305. Defendants' material misrepresentations and/or active concealment, suppression, and omissions constitute a continuing tort.

306. Through its package inserts and other public statements, Defendants continue to misrepresent the serious potential health risks and complications associated with use of Dupixent.

307. Defendants had a post-sale duty to timely warn physicians, including Plaintiff's healthcare providers, and consumers, such as Plaintiffs, about the potential risks and complications associated with use of Dupixent.

308. Defendants fraudulently and intentionally misrepresented the safety and efficacy of

Dupixent in their labeling, advertising, package inserts, promotional materials or other marketing resources and materials.

309. If Plaintiff's healthcare providers and Plaintiff had known the true facts concerning the risks of Dupixent use, in particular, the risk of CTCL, they would not have prescribed or used Dupixent and would have instead prescribed and used a safer alternative pharmaceutical drug or no drug at all.

310. Plaintiff and Plaintiff's healthcare providers' reliance upon Defendants' material misrepresentations were justified, among other reasons, because said misrepresentations and omissions were made by individuals and entities who were in a position of knowledge of the true facts concerning Dupixent, while Plaintiff and Plaintiff's healthcare providers were not in a position to know the true facts concerning Dupixent, and because Defendants overstated the benefits and safety of Dupixent, and concomitantly downplayed the risks of its use, including, but not limited to, an increased risk of CTCL, thereby inducing Plaintiff's healthcare providers to prescribe and Plaintiff to use Dupixent, in lieu of other safer alternatives, or no drug at all.

311. As a direct and proximate result of the dangerous and defective nature of Dupixent, Plaintiff suffered mycosis fungoides, and resulting pain and suffering, disability, disfigurement, mental anguish, loss of capacity for the enjoyment of life, expense of hospitalization, and medical care and treatment. These losses are permanent and the Plaintiff will continue to suffer the losses in the future.

FOURTH CAUSE OF ACTION NEGLIGENT MISREPRESENTATION

312. Plaintiff incorporates by reference herein each of the allegations heretofore set forth in this Complaint as though fully set forth herein.

313. Defendants' negligent material misrepresentations and omissions regarding the

safety and efficacy of Dupixent and of Dupixent's side effects, including that concerning an increased risk of CTCL, were communicated to Plaintiffs directly through promotional materials, advertising, package inserts, journal publications, and the product monograph with the intent that Plaintiff use Dupixent. The safety and efficacy of Dupixent was also negligently misrepresented to Plaintiff's healthcare providers with the intent that such misrepresentations would result in Dupixent being prescribed and administered to Plaintiff.

314. Defendants either knew or should have known that the material representations they were making regarding the safety, efficacy and side effects of Dupixent were false.

315. Defendants negligently made misrepresentations and/or actively concealed, suppressed, or omitted this material information with the intention and specific desire to induce consumers and the medical community, including Plaintiff and Plaintiff's healthcare providers, to use, prescribe and purchase Dupixent.

316. Defendants negligently knew or should have known that Plaintiff and/or Plaintiff's healthcare providers would rely upon such material misrepresentations and/or omissions in selecting Dupixent for the treatment of Plaintiff.

317. Plaintiff and Plaintiff's healthcare providers detrimentally relied on the negligent misrepresentations outlined below.

318. Defendants made these material misrepresentations and/or omissions and actively concealed adverse information at a time when they, their agents and/or their employees knew that Dupixent had certain defects, dangers and characteristics that differed from what had been represented to the medical community and the consuming public, including Plaintiff's healthcare providers and Plaintiff.

319. Defendants misrepresented critical aspects of Dupixent's development and safety

evaluation, including its carcinogenic risks and the scope of its clinical testing. As intended, these misrepresentations induced reliance of health care providers and drove prescriptions of Dupixent including Plaintiff's Dupixent prescription.

- a. Defendants publicly portray Dupixent as safe, thereby implying that they have adequately evaluated its carcinogenic potential, despite never conducted formal studies to assess mechanisms by which Dupixent may cause CTCL or accelerate its progression;
- b. Defendants misrepresented the results of premarket clinical trials and postmarketing observational studies evidencing a serious risk of CTCL with the use of Dupixent; and
- c. Defendants promoted Dupixent as a safe and effective treatment for patients lacking diagnostic confirmation of atopic dermatitis and other inflammatory conditions despite the fact that the Defendants had not properly studied Dupixent in that patient population.

320. Defendants partnered, funded, and leveraged influential “third parties” to misrepresent the safety and effectiveness of Dupixent and promote its use in an unrestricted manner inconsistent with its FDA-approved indications. Defendants’ negligent conduct, as outlined below, led to the treatment guidelines health care providers rely upon to prescribe Dupixent to patients, including Plaintiff.

- a. Through a 2017 “steering committee” funded by Defendants, which later published in *The Journal of Allergy and Clinical Immunology: In Practice*, Defendants falsely characterized Dupixent’s safety profile as being superior to existing therapies, including in terms of lymphoma risk, notwithstanding that it had not yet been used by any patients under real-world conditions;
- b. Defendants used this self-funded publication within their treatment guidelines, falsely recommending Dupixent as a first-line systemic treatment option in adults with moderate-to-severe AD despite the fact that: (1) Dupixent was not FDA-approved for any indication at the time the steering committee met or formulated its recommendations, and (2) Dupixent has never been approved by the FDA as a first-line systemic therapy for atopic dermatitis; and
- c. Defendants negligently represented this “steering committee” recommendations and manuscript were free from bias, despite the fact that 10 of the 11 authors of the guidelines article being retained by Defendants

as physician KOLs who had been paid a total of \$919,595 for consulting fees, food and beverage costs, education costs and travel and lodging expenses related to Dupixent.

321. Defendants, their agents and/or their employees, authored and funded misleading publications in the medical literature that contained false material facts and suppressed the truth regarding Dupixent's safety, efficiency, and association with CTCL, to which Defendants knew or should have known.

- a. Upon information and belief, and as discovery is expected to reveal, authors Yung-Tsu Cho and Chia-Yu Chu, acting at the direction of and in coordination with Defendants, published an August 2024 letter that sought to minimize the growing evidence linking Dupixent to CTCL, the precise injury suffered by Plaintiff, while failing to disclose Defendants' funding, involvement, and/or influence over the publication. This misleading communication was disseminated despite Defendants' knowledge of CTCL cases observed during the Dupixent clinical development program including at least three CTCL cases reported in the long-term extension trial.
- b. Despite there being 4 retrospective cohort studies, 2 disproportionality analyses and no fewer than 20 case reports and case series encompassing dozens of patients that had been published reporting an association between Dupixent and CTCL, on September 30, 2024, 10 employees of Defendants, Regeneron Pharmaceuticals, Inc. and Sanofi-Aventis U.S. LLC published a letter misrepresenting material scientific evidence concerning the rate of CTCL among the atopic dermatitis population and their knowledge of the existence of a safety signal for CTCL with Dupixent, omitting that their marketing targeted patients with less severe forms of atopic dermatitis – the opposite of the population Defendants asserted was likely to comprise the Dupixent treated group in the Hasan 2024 study;
- c. Defendants further perpetrated negligently misrepresentations and omissions by coordinating a series of publications authored by their retained and compensated key opinion leaders, including the Neubauer 2024 research letter, a contemporaneous TriNetX analysis, and a subsequent review article, that falsely minimized the Dupixent–CTCL safety signal, selectively omitted contrary data, misrepresented study findings and comparator groups, and concealed significant financial conflicts while urging clinicians not to allow CTCL evidence to influence their prescribing decisions;
- d. In September 2025, Defendants falsely represented that there is no evidence supporting an association between CTCL and Dupixent, while being aware

of at least 6 retrospective cohort studies, 5 disproportionality analyses and dozens of descriptive cohort studies, case reports, case series and systematic reviews. In addition, at that time, Defendants had already been notified by the FDA that the agency had identified a safety signal for CTCL with Dupixent;

- e. A published letter in June 2025 by Defendants' retained KOLs falsely suggested that heightened clinician "vigilance" led to over-reporting of CTCL with Dupixent, despite the well-known phenomenon of significant underreporting of postmarketing adverse events, and falsely asserting, without support, that no clear causal association had been identified;
- f. Defendants further advanced their negligently misrepresentations and omissions through a coordinated series of 2025 publications authored by their retained and heavily compensated KOLs, including Ian D. Pavord, Tiago Torres, and Adam Friedman, MD, which collectively downplayed the strength of emerging epidemiologic evidence linking Dupixent to CTCL, mischaracterized high-quality studies as "anecdotal," minimized the magnitude and seriousness of observed risks, urged clinicians not to let CTCL concerns deter prescribing, selectively emphasized Dupixent's purported benefits while concealing and misrepresenting causal mechanisms, and failed to disclose substantial financial conflicts despite publicly claiming none existed;
- g. In January 2026, through their retained KOLs, Defendants mischaracterized results of the Kridin 2026 study and Talasila 2026 research letter as reassuring evidence of Dupixent's safety by portraying the findings as reflecting a "decreased risk association," "partial amelioration" of lymphoma risk, or mere "diagnostic confounding and delayed recognition", despite conversely identifying an increased risk of CTCL with Dupixent in multiple populations, consistent with other studies reporting statistically significant increased risks of lymphoma among Dupixent-treated patients, while further providing no reliable basis to exclude a causal association with CTCL and mischaracterizing and/or omitting other available evidence relevant to the Dupixent-CTCL relationship.

322. Despite having a wealth of knowledge, since 2017, Defendants have at all times failed to disclose that the prescribing information for Dupixent is materially deficient. Each of the following failures constitute a negligent omission of safety information necessary for patients and prescribers, including Plaintiff, to be informed on the prescription and use of Dupixent:

- a. failure to include a BOXED WARNING advising patients and their healthcare providers of the risk of CTCL or any other form of cancer with Dupixent treatment;

- b. failure to include warnings in section 5, WARNINGS AND PRECAUTIONS, advising patients and their healthcare providers of the risk of CTCL or any other form of cancer with Dupixent treatment;
- c. failure to reference to CTCL in section 6, ADVERSE REACTIONS, advising patients and their healthcare providers that clinically significant CTCL adverse reactions have been reported with the use of Dupixent;
- d. failure to include a contraindication for use in patients with a history of CTCL or active CTCL;
- e. failure to include instructions directing healthcare providers to perform adequate testing of patients (through biopsy or other means) to confirm a clinical diagnosis of atopic dermatitis, particularly in equivocal cases, before commencing treatment with Dupixent;
- f. failure to include instructions directing healthcare providers to perform adequate testing of patients (through biopsy, cytometry or other means) to rule out the presence of subclinical or smoldering CTCL before commencing treatment with Dupixent;
- g. failure to include instructions directing healthcare providers to perform regular and continuous monitoring during treatment with Dupixent for the development of signs and symptoms of CTCL and repeat biopsies to detect changes in pathologies during treatment with Dupixent consistent with CTCL development; and
- h. failure to include instructions directing healthcare providers to discontinue use of Dupixent when a patient exhibits signs of non-responsiveness to treatment or worsening of disease, or otherwise develops signs and symptoms consistent with CTCL.

323. Additionally, Defendants affirmatively highlighted the absence of a boxed warning and the lack of any requirement for initial laboratory testing or ongoing laboratory monitoring as favorable safety attributes of Dupixent in their promotional and patient-facing materials. In doing so, Defendants mislead patients and prescribing physicians that Dupixent posed no serious or latent safety risks and had been adequately evaluated for long-term adverse outcomes.

324. To drive profits, Defendants intentionally misrepresented and omitted material information through a multifaceted marketing strategy, spanning direct-to-consumer advertising, disease-awareness initiatives, and other promotional mechanisms. The following marketing

strategies, executed by Defendants, their agents and/or their employees, were relied on by Plaintiff and her prescribing physician:

- a. Defendants misrepresented Dupixent's safety and effectiveness through provocative direct-to-consumer advertising that used glossy imagery and slogans such as "Du-More," "Better Days," and "This is Better" to falsely imply that patients would achieve materially happier, healthier, and more fulfilling lives by using Dupixent;
- b. Defendants further executed negligently misrepresentations and omissions by deploying the Dupixent MyWay and MyWay Ambassador programs, related smartphone applications, and a network of unbranded disease-state awareness campaigns and websites, including UnderstandAD.com, EczemaExposed.com, ThinkCOPDInflammation.com, and LiveWithCOPD.com, to create the false impression of neutral educational resources while in fact steering patients to view their symptoms as serious and uncontrolled, encouraging "productive" doctor conversations designed to yield Dupixent prescriptions, and seeding demand for Dupixent even before FDA approval, all while concealing that these materials were crafted and controlled by Defendants to promote Dupixent's use;
- c. Defendants fraudulently misrepresented diagnostic tools, including the Atopic Dermatitis Control Tool (ADCT) and related pediatric itch scales neutral clinical instruments, when in fact their high sensitivity artificially inflated atopic dermatitis diagnoses and artificially steered patients and physicians toward Dupixent treatment; and
- d. Through continuing medical education programs and their ADVENT global medical education platform, Defendants misrepresented promotional materials as unbiased scientific education, concealing their true marketing intent to increase Dupixent prescribing.

325. At no point did Defendants acknowledge the merits of the findings produced through independent research or take initiative to commit to conducting necessary studies to better understand the pathological mechanisms of Dupixent-induced CTCL or discover why Dupixent triggers rapid and severe progression of subclinical CTCL, instead Defendants misrepresented material facts to the public by funding and influence.

326. As further detailed above, Defendants engaged in the following negligent practices designed to induce healthcare providers and consumers to prescribe and use Dupixent, which were

relied on by Plaintiff's prescribing physician:

- a. inducing healthcare providers to switch their patients from other medications to Dupixent despite lacking the necessary evidence to demonstrate that Dupixent is safe and effective in this population and despite affirmative evidence that the drug is not safe in this patient population;
- b. inducing healthcare providers, through marketing efforts, to prescribe Dupixent to their patients without first obtaining diagnostic confirmation of an underlying indicated health condition;
- c. engaging in marketing efforts, such as disease state awareness campaigns and branded advertising campaigns, seeking to induce patients to obtain a diagnosis for an indicated health condition and then specifically request a prescription of Dupixent from their healthcare provider when its use is not necessary; and
- d. intentionally misleading healthcare providers and the general public by making superiority claims for Dupixent as compared to other medications despite possessing the knowledge that these claims are false.

327. Defendants' misrepresentations were widely disseminated to healthcare providers through CME programs, clinical guidelines, KOL publications, and disease awareness campaigns, and to consumers via direct-to-consumer advertising, MyWay materials, and unbranded disease-state websites. These misrepresentations reached Plaintiff and Plaintiff's treating providers and directly influenced the decision to prescribe and use Dupixent.

328. Defendants' material misrepresentations and/or active concealment, suppression, and omissions were perpetuated directly and/or indirectly by Defendants, their sales representatives, employees, distributors, agents and/or detail persons, through databases, printouts, monographs, product labeling and other information drafted, prepared, marketed, sold, and supplied by Defendants, their sales representatives, employees, distributors, agents and/or detail persons.

329. Defendants' material misrepresentations and/or active concealment, suppression, and omissions constitute a continuing tort.

330. Through its package inserts and other public statements, Defendants continue to misrepresent the serious potential health risks and complications associated with use of Dupixent.

331. Defendants had a post-sale duty to timely warn physicians including Plaintiff's healthcare providers, and consumers, such as Plaintiffs, about the potential risks and complications associated with use of Dupixent.

332. Defendants negligently misrepresented the safety and efficacy of Dupixent in their labeling, advertising, package inserts, promotional materials or other marketing resources and materials.

333. If Plaintiff's healthcare providers and Plaintiff had known the true facts concerning the risks of Dupixent use, in particular, the risk of CTCL, they would not have prescribed or used Dupixent and would have instead prescribed and used a safer alternative pharmaceutical drug or no drug at all.

334. Plaintiff and Plaintiff's healthcare providers' reliance upon Defendants' material misrepresentations were justified, among other reasons, because said misrepresentations and omissions were made by individuals and entities who were in a position of knowledge of the true facts concerning Dupixent, while Plaintiff and Plaintiff's healthcare providers were not in a position to know the true facts concerning Dupixent, and because Defendants overstated the benefits and safety of Dupixent, and concomitantly downplayed the risks of its use, including, but not limited to, an increased risk of CTCL, thereby inducing Plaintiff's healthcare providers to prescribe and Plaintiff to use Dupixent, in lieu of other safer alternatives, or no drug at all.

335. As a direct and proximate result of the dangerous and defective nature of Dupixent, Plaintiff suffered mycosis fungoides, and resulting pain and suffering, disability, disfigurement, mental anguish, loss of capacity for the enjoyment of life, expense of hospitalization, and medical

care and treatment. These losses are permanent and the Plaintiff will continue to suffer the losses in the future.

FIFTH CAUSE OF ACTION
PUNITIVE DAMAGES

336. Plaintiff incorporates by reference herein each of the allegations heretofore set forth in this Complaint as though fully set forth herein.

337. As discussed herein, Defendants have intentionally misrepresented the clinical trial and postmarketing safety data for Dupixent to FDA, healthcare providers, and the general public in order to mask the true risk of CTCL related to Dupixent use. Those misrepresentations continue to the present.

338. Defendants have engaged in marketing efforts seeking to induce healthcare providers to switch their patients from other medications to Dupixent despite lacking the necessary evidence to demonstrate that Dupixent is safe and effective in this population and despite affirmative evidence that the drug is not safe in this patient population.

339. Defendants have engaged in marketing efforts seeking to induce healthcare providers to prescribe Dupixent to their patients without first obtaining diagnostic confirmation of an underlying indicated health condition.

340. Defendants have engaged in marketing efforts seeking to induce patients to specifically request a prescription of Dupixent from their healthcare provider when its use is not necessary.

341. Defendants have intentionally misled healthcare providers and the general public in making superiority claims for Dupixent as compared to other medications despite possessing the knowledge that these claims are false.

342. Defendants have intentionally failed to properly warn healthcare providers about

the true risk of CTCL related to Dupixent use despite possessing knowledge that Dupixent causes these serious adverse events.

343. Defendants' actions were willful and malicious in that Defendants' conduct was carried on with a conscious disregard for the safety and rights of Plaintiff. Defendants' unconscionable conduct thereby warrants an assessment of exemplary and punitive damages against Defendants in an amount appropriate to punish Defendants, and deter similar conduct in the future.

344. As a direct and proximate result of the dangerous and defective nature of Dupixent, Plaintiff suffered mycosis fungoides, and resulting pain and suffering, disability, disfigurement, mental anguish, loss of capacity for the enjoyment of life, expense of hospitalization, and medical care and treatment. These losses are permanent and the Plaintiff will continue to suffer the losses in the future.

PRAYER FOR RELIEF

WHEREFORE, Plaintiff prays for judgment against Defendants, as follows:

- a. For general damages in an amount to be proven at the time of trial;
- b. For special damages in an amount to be proven at the time of trial;
- c. For lost wages and/or lost earning capacity;
- d. For exemplary and punitive damages in an amount to be proven at trial, and sufficient to punish Defendants or to deter Defendants and others from repeating the injurious conduct alleged herein;
- e. For pre-judgment and post-judgment interest on the above general and special damages;
- f. For costs of this suit and attorneys' fees; and

g. All other relief that this Court deems necessary, proper, and just.

DEMAND FOR JURY TRIAL

Plaintiff demands a trial by jury on all issues.

Respectfully submitted,

Dated: August 3, 2026

/s/ Matthew R. Fogg

Matthew R. Fogg, Esq. DE Bar #4254

MORRIS JAMES LLP

3205 Avenue North Blvd., Ste. 100

Wilmington, DE 19803

mfogg@morrisjames.com

(302) 655-8808 (Telephone)

/s/ Keith E. Donovan

Keith E. Donovan, Esq. DE Bar #3044

MORRIS JAMES LLP

3205 Avenue North Blvd., Ste. 100

Wilmington, DE 19803

kdonovan@morrisjames.com

(302) 655-8808 (Telephone)

/s/ Brandon L. Bogle

Brandon L. Bogle, Esq. FL Bar #52624

**Levin, Papantonio, Proctor, Buchanan,
O'Brien, Barr & Mougey, P.A.**

316 South Baylen Street, Suite 600

Pensacola, Florida 32502

bbogle@levinlaw.com

(850) 435-7043 (Telephone)

/s/ Madison A. Mougey

Madison A. Mougey, Esq. FL Bar # 1071135

**Levin, Papantonio, Proctor, Buchanan,
O'Brien, Barr & Mougey, P.A.**

316 South Baylen Street, Suite 600

Pensacola, Florida 32502

mmougey@levinlaw.com

(850) 435-7043 (Telephone)

Attorneys for Plaintiff

CIVIL COVER SHEET

The JS 44 civil cover sheet and the information contained herein neither replace nor supplement the filing and service of pleadings or other papers as required by law, except as provided by local rules of court. This form, approved by the Judicial Conference of the United States in September 1974, is required for the use of the Clerk of Court for the purpose of initiating the civil docket sheet. (SEE INSTRUCTIONS ON NEXT PAGE OF THIS FORM.)

I. (a) PLAINTIFFS

Anne Gibson

(b) County of Residence of First Listed Plaintiff Grand County, Colorado
(EXCEPT IN U.S. PLAINTIFF CASES)

(c) Attorneys (Firm Name, Address, and Telephone Number)

Levin, Papantonio, Proctor, Buchanan, O'Brien, Barr & Mougey, P.A.
316 South Baylen Street, Suite 600, Pensacola, Florida 32502
(850) 435-7043 (Telephone)

DEFENDANTS

Sanofi-Aventis U.S. LLC, Genzyme Corporation, and Regeneron Pharmaceuticals, Inc

County of Residence of First Listed Defendant Somerset County
(IN U.S. PLAINTIFF CASES ONLY)

NOTE: IN LAND CONDEMNATION CASES, USE THE LOCATION OF THE TRACT OF LAND INVOLVED.

Attorneys (If Known)

II. BASIS OF JURISDICTION (Place an "X" in One Box Only)

- ☐ 1 U.S. Government Plaintiff ☐ 3 Federal Question (U.S. Government Not a Party)
- ☐ 2 U.S. Government Defendant ☒ 4 Diversity (Indicate Citizenship of Parties in Item III)

III. CITIZENSHIP OF PRINCIPAL PARTIES (Place an "X" in One Box for Plaintiff and One Box for Defendant)

- | | PTF | DEF | | PTF | DEF |
|---|---------------------------------------|----------------------------|---|----------------------------|---------------------------------------|
| Citizen of This State | ☐ 1 | ☐ 1 | Incorporated or Principal Place of Business In This State | ☐ 4 | ☒ 4 |
| Citizen of Another State | ☒ 2 | ☐ 2 | Incorporated and Principal Place of Business In Another State | ☐ 5 | ☐ 5 |
| Citizen or Subject of a Foreign Country | ☐ 3 | ☐ 3 | Foreign Nation | ☐ 6 | ☐ 6 |

IV. NATURE OF SUIT (Place an "X" in One Box Only)Click here for: [Nature of Suit Code Descriptions.](#)

CONTRACT	TORTS	FORFEITURE/PENALTY	BANKRUPTCY	OTHER STATUTES	
☐ 110 Insurance ☐ 120 Marine ☐ 130 Miller Act ☐ 140 Negotiable Instrument ☐ 150 Recovery of Overpayment & Enforcement of Judgment ☐ 151 Medicare Act ☐ 152 Recovery of Defaulted Student Loans (Excludes Veterans) ☐ 153 Recovery of Overpayment of Veteran's Benefits ☐ 160 Stockholders' Suits ☐ 190 Other Contract ☐ 195 Contract Product Liability ☐ 196 Franchise	PERSONAL INJURY ☐ 310 Airplane ☐ 315 Airplane Product Liability ☐ 320 Assault, Libel & Slander ☐ 330 Federal Employers' Liability ☐ 340 Marine ☐ 345 Marine Product Liability ☐ 350 Motor Vehicle ☐ 355 Motor Vehicle Product Liability ☐ 360 Other Personal Injury ☐ 362 Personal Injury - Medical Malpractice	PERSONAL INJURY ☐ 365 Personal Injury - Product Liability ☒ 367 Health Care/Pharmaceutical Personal Injury Product Liability ☐ 368 Asbestos Personal Injury Product Liability PERSONAL PROPERTY ☐ 370 Other Fraud ☐ 371 Truth in Lending ☐ 380 Other Personal Property Damage ☐ 385 Property Damage Product Liability	☐ 625 Drug Related Seizure of Property 21 USC 881 ☐ 690 Other LABOR ☐ 710 Fair Labor Standards Act ☐ 720 Labor/Management Relations ☐ 740 Railway Labor Act ☐ 751 Family and Medical Leave Act ☐ 790 Other Labor Litigation ☐ 791 Employee Retirement Income Security Act IMMIGRATION ☐ 462 Naturalization Application ☐ 465 Other Immigration Actions	☐ 422 Appeal 28 USC 158 ☐ 423 Withdrawal 28 USC 157 INTELLECTUAL PROPERTY RIGHTS ☐ 820 Copyrights ☐ 830 Patent ☐ 835 Patent - Abbreviated New Drug Application ☐ 840 Trademark ☐ 880 Defend Trade Secrets Act of 2016 SOCIAL SECURITY ☐ 861 HIA (1395ff) ☐ 862 Black Lung (923) ☐ 863 DIWC/DIWW (405(g)) ☐ 864 SSID Title XVI ☐ 865 RSI (405(g)) FEDERAL TAX SUITS ☐ 870 Taxes (U.S. Plaintiff or Defendant) ☐ 871 IRS—Third Party 26 USC 7609	☐ 375 False Claims Act ☐ 376 Qui Tam (31 USC 3729(a)) ☐ 400 State Reapportionment ☐ 410 Antitrust ☐ 430 Banks and Banking ☐ 450 Commerce ☐ 460 Deportation ☐ 470 Racketeer Influenced and Corrupt Organizations ☐ 480 Consumer Credit (15 USC 1681 or 1692) ☐ 485 Telephone Consumer Protection Act ☐ 490 Cable/Sat TV ☐ 850 Securities/Commodities/Exchange ☐ 890 Other Statutory Actions ☐ 891 Agricultural Acts ☐ 893 Environmental Matters ☐ 895 Freedom of Information Act ☐ 896 Arbitration ☐ 899 Administrative Procedure Act/Review or Appeal of Agency Decision ☐ 950 Constitutionality of State Statutes
REAL PROPERTY ☐ 210 Land Condemnation ☐ 220 Foreclosure ☐ 230 Rent Lease & Ejectment ☐ 240 Torts to Land ☐ 245 Tort Product Liability ☐ 290 All Other Real Property	CIVIL RIGHTS ☐ 440 Other Civil Rights ☐ 441 Voting ☐ 442 Employment ☐ 443 Housing/Accommodations ☐ 445 Amer. w/Disabilities - Employment ☐ 446 Amer. w/Disabilities - Other ☐ 448 Education	PRISONER PETITIONS Habeas Corpus: ☐ 463 Alien Detainee ☐ 510 Motions to Vacate Sentence ☐ 530 General ☐ 535 Death Penalty Other: ☐ 540 Mandamus & Other ☐ 550 Civil Rights ☐ 555 Prison Condition ☐ 560 Civil Detainee - Conditions of Confinement			

V. ORIGIN (Place an "X" in One Box Only)

- ☐ 1 Original Proceeding ☐ 2 Removed from State Court ☐ 3 Remanded from Appellate Court ☐ 4 Reinstated or Reopened ☐ 5 Transferred from Another District (specify) ☐ 6 Multidistrict Litigation - Transfer ☒ 8 Multidistrict Litigation - Direct File

VI. CAUSE OF ACTION

Cite the U.S. Civil Statute under which you are filing (Do not cite jurisdictional statutes unless diversity):

28 U.S.C. 1332; 28 USC 1391

Brief description of cause:

VII. REQUESTED IN COMPLAINT:

☐ CHECK IF THIS IS A CLASS ACTION UNDER RULE 23, F.R.Cv.P.

DEMAND \$

CHECK YES only if demanded in complaint:

JURY DEMAND: ☒ Yes ☐ No**VIII. RELATED CASE(S) IF ANY**

(See instructions):

JUDGE ZAHID N. QURAISHIDOCKET NUMBER 3:26-md-03180-ZNQ-JBD

DATE

8/3/2026

SIGNATURE OF ATTORNEY OF RECORD

/s/ Brandon L. Bogle

FOR OFFICE USE ONLY

RECEIPT # _____ AMOUNT _____ APPLYING IFP _____ JUDGE _____ MAG. JUDGE _____